

Astroglial changes in the zona incerta in response to motor cortex stimulation in a rat model of chronic neuropathy

Myeounghoon Cha¹, Kyung Hee Lee², Bae Hwan Lee^{1,3*}

¹ Department of Physiology, Yonsei University College of Medicine, Seoul 03722, Republic of Korea

² Department of Dental Hygiene, Division of Health Science, Dongseo University, Busan 47011, Republic of Korea

³ Brain Korea 21 PLUS Project for Medical Science, Brain Research Institute, Epilepsy Research Institute, Yonsei University College of Medicine, Seoul 03722, Republic of Korea

* Corresponding Author

Bae Hwan Lee, PhD

Department of Physiology, Yonsei University College of Medicine


50-1, Yonsei-ro, Seodaemun-gu

Seoul 03722, Korea

Tel.: +82.2.2228.1711


Fax: +82.2.393.0203

E-mail address: bhlee@yuhs.ac




Supplementary Figure S1. Full-length blots of western blotting shown in Figure 2E. Punctured the ZI tissues were subjected to western blotting with GFAP or NeuN antibodies. Anti- β -actin antibody was used as a loading control. The cropped parts indicate by the gray dashed lines are shown in Figure 2E

A



B



Supplementary Figure S2. Full-length blots of western blotting shown in Figure 5A and B. Punctured the M1 and ZI tissues were subjected to western blotting with MAP2, PSD95, and Synapsin antibodies. Anti-GAPDH antibody was used as a loading control. The cropped parts indicate by the gray dashed lines are shown in Figure 5A and B.