

Supporting Information for

**Synthesis of Conductive Carbon Aerogels
Decorated with β -Tricalcium Phosphate Nanocrystallites**


Atakan Tevlek¹, Abdulraheem M. N. Atya¹, Muhannad Almemar¹,

Memed Duman², Dincer Gokcen³, Alexey Y. Ganin⁴,

Humphrey H. P. Yiu⁵ and Halil M. Aydin^{1,6}

<i>Index</i>	<i>Page</i>
SI-1	2
SI-2	2
SI-3	3
SI-4	3
SI-5	4
SI-6	4
SI-7	5
SI-8	5
SI-9	6


SI-1. Schematic representation of the set-up for 4-probe measurement system.



SI-2. Micro-CT data of a representative β -TCP incorporated (Sample 850 °C 1XTCP) carbon aerogel structure.

Parameter	Value
Total Porosity	90.80 %
Open Porosity	90.75 %
Surface Area	795.14 mm ² /sample
Surface/Volume Ratio	16.79 1/mm
Connection Density	1.49 1/mm ³


SI-3: SEM images shows the β -TCP nanocrystallite sizes on microbelts (Sample 1100 °C 1XTCP):



Particle Size calculated by using Image J Software (1.52a USA)

Mean	74.76 nm
S.D.	30.17 nm
Minimum	30.27 nm
Maximum	199.33 nm


SI-4: TEM images of the constructs: carbon aerogels without (left) (Sample 850 °C - 0XTCP) and with (right) β -TCP (850 °C - 1XTCP):




SI-5: EDX atomic percentages (Sample 1100 °C 1XTCP).

Element	Atomic %
Calcium	2.79
Phosphorus	1.85
Oxygen	6.15
<i>Ca/P Ratio</i>	<i>1.5081</i>

SI-6: XRD diffractogram of β -TCP particles used in the synthesis step.




SI-7: XRD diffractogram of a reference β -TCP from the library.



(Source: <http://icsd.cds.rsc.org/search/basic.xhtml>)

SI-8. FTIR spectra of pure β -TCP particles and carbon aerogels (C.A.) with (1XTCP) and without (0XTCP) ceramic phase, both pyrolyzed at 850 °C.



The FTIR spectra of the prepared aerogels were analyzed. The broad band at around 3200 cm⁻¹ (stretching vibrations) and the band at around 1425 cm⁻¹ (bending vibrations) were due to the existence of -CH₃, -CH₂-, =CH- groups in the structure of pyrolysis products without β -TCP. For the case of β -TCP containing pulp pyrolysis, the signals representing C-H bonds completely disappeared due to the existence of three σ and one π bonds between only carbon atoms. Additionally, broad and weak C=C stretching vibrations at around 1565 cm⁻¹ and weak bending vibrations at around 725 cm⁻¹ also observed in the structure of the pyrolyzed pulp even though it contains three-fold higher amounts of β -TCP than the paper. Therefore, the broad band around 1000 cm⁻¹ and the peaks at 600 cm⁻¹ and 540 cm⁻¹ shows the existence of β -TCP, namely symmetric stretching mode ν_1 , vibrational modes ν_4 and ν_3 , respectively, in addition to the interfered IR signals of carbon material formed in the structure of the same pyrolyzed sample.

SI-9. Mean size values of the ordered domains of β -TCP ceramic nanocrystallites on the carbon aerogel fibers by XRD.

Sample	Mean FWHM	Calculated Crystallite Size (nm)
850 °C 1/2XTCP	0.265	35.01
850 °C 1XTCP	0.159	58.38
850 °C 2XTCP	0.219	42.40
1100 °C 1/2XTCP	0.157	59.75
1100 °C 1XTCP	0.158	58.75
1100 °C 2XTCP	0.151	61.51

*Scherer Equation ($\tau = K\lambda / \beta \cos\theta$) was used in the calculations. Shape factor K was taken as 0.9 while λ was taken as 0,15406 nm. β , the broadening at the FWHM intensity (radians) and Bragg angle (θ) was given in the table. The highest and characteristic three Bragg angle values were selected. 3 Bragg angles used for each diffractogram (e.g. 2Tetha=31.11°, 28.02°, 34.5°).