

RXR EXPRESSION IN MARINE GASTROPODS WITH DIFFERENT SENSITIVITY TO IMPOSEX DEVELOPMENT

Sebastián Giulianelli^{1,2}, Mónica A. Primost^{1,3}, Claudia Lanari² and Gregorio Bigatti^{1,4,5}

¹Instituto de Biología de Organismos Marinos. IBIOMAR-CCT CENPAT-CONICET. Puerto Madryn, Argentina.

²Instituto de Biología y Medicina Experimental. IBYME-CONICET. Buenos Aires, Argentina.

³Grupo de Investigación y Desarrollo Tecnológico en Acuicultura y Pesca (GIDTAP). Universidad Tecnológica Nacional. Facultad Regional Chubut (UTN-FRCh), Argentina.

⁴Universidad Nacional de la Patagonia San Juan Bosco. Puerto Madryn, Argentina.

⁵Universidad Espíritu Santo, Ecuador.

Supplementary Figure S1:




Figure S1. Study area and location of the sampled sites. The map was sourced from Google Earth Pro (v7.3) and modified in Adobe Illustrator (v16.0.0). CA: Cerro Avanzado beach, LPBH: Luis Piedra Buena Harbor.

Supplementary Figure S2:





Figure S2. Immunohistochemical staining for RXR. (a and b) Epidermal region of male penis from *B. globulosus* and *T. geversianus*, respectively. Bar: 50 μ m. (c and d) Gonads of *B. globulosus* and *T. geversianus* females, respectively. Bar: 50 μ m. (e and f) Digestive glands of *B. globulosus* and *T. geversianus* females, respectively. Bar: 50 μ m.

Supplementary Figure S3: Full-length gels from Figure 2



Supplementary Figure S4: Full-length gels from Figure 3

