

SUPPLEMENTAL MATERIAL

Full title: Epidemiology and Treatment of Atrial Fibrillation in Patients with Type 2 Diabetes in the UK, 2001-2016.

Hassan Alwafi, MD^{1,2}, Ian CK Wong, PhD^{1,3}, Amitava Banerjee, DPhil^{4,5}, Pajaree Mongkhon, PhD^{6,7}, Cate Whittlesea, PhD¹, Abdallah Y Naser, PhD⁸, Wallis C.Y. Lau, PhD^{1,3}, Li Wei, PhD¹

¹Research Department of Practice and Policy, School of Pharmacy, University College London, London, United Kingdom

²Faculty of Medicine, Umm Al Qura University, Mecca, Saudi Arabia

³Centre for Safe Medication Practice and Research, Department of Pharmacology and Pharmacy, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong

⁴Institute of Health Informatics, University College London, London, UK

⁵Barts Health NHS Trust, London, UK

⁶Division of Pharmacy Practice, Department of Pharmaceutical Care, School of Pharmaceutical Sciences, University of Phayao, Phayao, Thailand

⁷Pharmacoepidemiology and Statistics Research Center (PESRC), Faculty of Pharmacy, Chiang Mai University, Chiang Mai, Thailand

⁸Faculty of Pharmacy, Isra University, Amman, Jordan

Corresponding author

Professor Li Wei
UCL School of Pharmacy
29-39 Brunswick Square
London, WC1N 1AX
T: 020 7874 1275
E: l.wei@ucl.ac.uk

Tables (1)

Table S1. Interrupted time series analysis model on the changes before and after the introduction of DOACs in OAC prescribing

Table S1: Interrupted time series analysis model on the changes before and after the introduction of DOACs in OAC prescribing.

Variable	IRR (95%CI)	Standard Error	P-value
Change in level (immediate effect)	1.00 (0.99-1.01)	0.056	0.299
Change in trend post intervention	1.02 (1.01-1.03)	0.004	<.0001

Abbreviations: CI, confidence interval; IRR, incidence rate ratio;