


Recombinant *Plasmodium vivax* Circumsporozoite Surface Protein allelic variants: antibody recognition by individuals from three communities in the Brazilian Amazon

Isabela Ferreira Soares^{#1}, César López-Camacho^{#2}, Rodrigo Nunes Rodrigues-da-Silva^{#3}, Ada da Silva Matos¹, Barbara de Oliveira Baptista⁴, Paulo Renato Rivas Totino⁴, Rodrigo Medeiros de Souza⁵, Kate Harrison², Alba Marina Gimenez², Elisângela Oliveira de Freitas², Young Chan Kim², Joseli Oliveira-Ferreira¹, Cláudio Tadeu Daniel-Ribeiro^{4,6}, Arturo Reyes-Sandoval², Lilian Rose Pratt-Riccio⁴, Josué da Costa Lima-Junior¹



Supplementary figure 1: Frequencies and RI's of IgG responders and subclasses separated by infected and non-infected individuals. White bars and white circles represent infected individuals while dark bars and dark circles represent non-infected individuals. Figures (a) and (b) compare frequencies and RI's of infected and non-infected individuals responding to each recombinant protein. Figures (c,d,e,f,g,h,i and j) compare frequencies and RI's of infected and non-infected individuals' IgG subclasses to each recombinant protein. Significant differences were indicated by *. (*) $p < 0.05$; (**) $p < 0.005$; *** $p < 0.0005$.