

Extracellular vesicles from amyloid- β exposed cell cultures induce severe dysfunction in cortical neurons

Chiara Beretta¹, Elisabeth Nikitidou¹, Linn Streubel-Gallasch¹, Martin Ingelsson¹, Dag Sehlin¹
and Anna Erlandsson¹

¹ Department of Public Health & Caring Sciences, Molecular Geriatrics, Rudbeck Laboratory, Uppsala University, Uppsala, Sweden

Corresponding author

Anna Erlandsson

Department of Public Health and Caring Sciences / Molecular Geriatrics,
Rudbeck Laboratory, Uppsala University

SE-751 85 Uppsala, Sweden

E-mail: anna.erlandsson@pubcare.uu.se

Co-authors

Chiara Beretta

Department of Public Health and Caring Sciences / Molecular Geriatrics,
Rudbeck Laboratory, Uppsala University

SE-751 85 Uppsala, Sweden

E-mail: chiara.beretta@pubcare.uu.se

Elisabeth Nikitidou

Department of Public Health and Caring Sciences / Molecular Geriatrics,

Rudbeck Laboratory, Uppsala University

SE-751 85 Uppsala, Sweden

E-mail: *elisabeth.nikitidou@telia.com*

Linn Streubel-Gallasch

Department of Public Health and Caring Sciences / Molecular Geriatrics,

Rudbeck Laboratory, Uppsala University

SE-751 85 Uppsala, Sweden

E-mail: *linn.streubel-gallasch@pubcare.uu.se*

Martin Ingelsson

Department of Public Health and Caring Sciences / Molecular Geriatrics,

Rudbeck Laboratory, Uppsala University

SE-751 85 Uppsala, Sweden

E-mail: *martin.ingelsson@pubcare.uu.se*


Dag Sehlin

Department of Public Health and Caring Sciences / Molecular Geriatrics,

Rudbeck Laboratory, Uppsala University


SE-751 85 Uppsala, Sweden

E-mail: *dag.sehlin@pubcare.uu.se*



Supplementary Figure 1.

Unprocessed, full membrane images of the Western blot analysis of synaptophysin and β -actin shown in Figure 5.



Supplementary Figure 2.

A zoomed out image of the stained cell culture shown in Figure 7 c, illustrating that the membranes of the neurons are positive for the cholesterol-binding dye filipin III. Scale bar: c=20 μ m.