

Complete Intraureteral Stent Placement Relieves Daytime Urinary Frequency Compared with Conventional Placement in Patients with an Indwelling Ureteral Stent: Post-hoc Analysis of a Randomized, Controlled Trial

Tomoaki Matsuzaki^{1†}, Takashi Yoshida^{1,2†}, Takashi Murota^{1,3}, Kazuyoshi Nakao¹, Makoto Taguchi¹, Hidefumi Kinoshita¹, and Tadashi Matsuda¹

¹Department of Urology and Andrology, Kansai Medical University, Osaka, Japan

²Department of Urology and Andrology, Kori Hospital, Kansai Medical University, Osaka, Japan

³Department of Urology and Andrology, General Medical Center, Kansai Medical University, Osaka, Japan


†These authors contributed equally to the work

Corresponding Author: Takashi Yoshida, MD

Department of Urology and Andrology, Kansai Medical University, 2-3-1 Shinmachi, Hirakata, Osaka 573-1191, Japan

Tel: +81-72-804-0101; fax: +81-72-804-2089

E-mail: yoshidtk@takii.kmu.ac.jp



Supplementary Figure 1. Study design of the post-hoc analysis. *Lost to follow-up (n=1), withdrew consent (n=2), and insufficient clinical data (n=1). **Lost to follow-up (n=1). I-PSS, International Prostate Symptom Score; and OABSS, Overactive Bladder Symptom Score.