

Renal Function is Associated with Blood Neurofilament Light Chain Level in Older Adults

Shoshin Akamine^{1,2}, Noriko Marutani^{2,3}, Daisuke Kanayama^{1,2,4}, Shiho Gotoh⁴, Riki Maruyama^{1,2}, Kanta Yanagida^{1,2}, Yukako Sakagami^{1,2}, Kohji Mori⁴, Hiroyoshi Adachi^{1,2,4}, Junji Kozawa^{5,6}, Norikazu Maeda^{5,7}, Michio Otsuki⁵, Takaaki Matsuoka⁵, Hiromi Iwahashi^{5,6,8}, Iichiro Shimomura⁵, Manabu Ikeda⁴, Takashi Kudo^{1,2,4}

Department of Mental Health Promotion, Osaka University Graduate School of Medicine, Suita, Osaka, Japan.

Health and Counseling Center, Osaka University, Toyonaka, Osaka, Japan.

Department of Psychological Health Promotion, Osaka University Graduate School of Medicine, Suita, Osaka, Japan.

Department of Psychiatry, Osaka University Graduate School of Medicine, Suita, Osaka, Japan.

Department of Metabolic Medicine, Osaka University Graduate School of Medicine, Suita, Osaka, Japan.

Department of Diabetes Care Medicine, Osaka University Graduate School of Medicine, Suita, Osaka, Japan.


Department of Metabolism and Atherosclerosis, Osaka University Graduate School of Medicine, Suita, Osaka, Japan.

Department of Internal Medicine, Toyonaka Municipal Hospital, Osaka, Japan.


Supplementary Information

Supplementary Figure 1. Scatterplot between blood NfL level and age (a), sex (b) in the HC group. NfL: neurofilament light chain.

(a)




(b)




Supplementary Figure 2. Scatterplot between blood NfL level and age (a), sex (b) in the DM group. NfL: neurofilament light chain.

(a)




(b)



Supplementary Figure 3. Q-Q plot before and log-transformation of serum creatinine level and blood neurofilament level in the (a) HC group and (b) DM group.

(a)



(b)

