

Antimicrobial peptides in patients with anorexia nervosa: comparison with healthy controls and the impact of weight gain

Marie-Christin Bendix, Michael Stephan, Mariel Nöhre, Wally Wunsch-Leiteritz, Hagen Schmidt, Gisa Tiegs, Jürgen Harder, Martina de Zwaan

Table 1 supplement. Pearson's correlations between baseline AMP concentrations in patients with AN with age, illness duration, current BMI, lowest and highest lifetime BMI.

	Age	Illness duration	Current BMI	Lowest BMI	Highest BMI
Psoriasin (ng/ml) forehead					
r	-.095	-.189	.169	.085	.118
p	.599	.291	.348	.638	.513
Psoriasin (ng/ml) forearm proximal					
r	.266	.311	.079	.008	.110
p	.134	.078	.661	.964	.542
Psoriasin (ng/ml) forearm distal					
r	.265	.100	-.202	-.222	.021
p	.136	.579	.259	.215	.906
RNase 7 (ng/ml) forehead					
r	.028	-.144	.169	.085	.141
p	.875	.422	.348	.638	.434
RNase 7 (ng/ml) forearm proximal					
r	-.056	-.091	.212	.211	.044
p	.756	.614	.236	.238	.808
RNase 7 (ng/ml) forearm distal					
r	-.045	-.062	.271	.137	.138
p	.805	.730	.128	.447	.444

r = correlation coefficient