

Supplementary figure 1. Correlation between the expression of these candidate cirRNAs in normal and tumour samples

Title:

Genome-wide identification and characterization of circular RNA in resected hepatocellular carcinoma and background liver tissue

Authors:

Yuki Sunagawa, MD[†]; Suguru Yamada^{*}, MD, PhD; Fuminori Sonohara, MD, PhD[†]; Keisuke Kurimoto, MD, PhD; Nobutake Tanaka, MD, PhD; Yunosuke Suzuki, MD; Yoshikuni Inokawa, MD, PhD; Hideki Takami, MD, PhD; Masamichi Hayashi, MD, PhD; Mitsuro Kanda, MD, PhD; Chie Tanaka, MD, PhD; Goro Nakayama, MD, PhD; Masahiko Koike, MD, PhD; and Yasuhiro Koderu, MD, PhD

[†] These authors contributed equally to this work.

Affiliations:

Department of Gastroenterological Surgery, Nagoya University Graduate School of Medicine, Nagoya, Japan

*Corresponding author:

Suguru Yamada, MD, PhD

Department of Gastroenterological Surgery, Nagoya University Graduate School of Medicine, 65, Tsurumai-cho, Showa-ku, Nagoya, 466-8550, Japan

Tel: +81-52-744-2245; Fax: +81-52-744-2255; Email: suguru@med.nagoya-u.ac.jp

Legend:

The figure is prepared using R software (version 3.5.3, <http://www.r-project.org/>) [43].

Supplementary figure 1. Correlation between the expression of these candidate cirRNAs in normal and tumour samples

