

**Microvesicles released from pneumolysin-stimulated lung epithelial cells carry
mitochondrial cargo and suppress neutrophil oxidative burst**

Letsiou E.^{1,2*}, Teixeira Alves LG.¹, Fatykhova D.¹, Felten M.¹, Mitchell TJ.³, Mueller-Redetzky H.¹, Hocke AC.^{1,4}, Witzernath M.^{1,4}


¹Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Division of Pulmonary Inflammation, and Department of Infectious Diseases and Respiratory Medicine, Charitéplatz 1, 10117 Berlin, Germany.

²Division of Pulmonary, Critical Care, Sleep, and Allergy, University of Illinois at Chicago, 60612 Chicago, Illinois, USA.


³Institute of Microbiology and Infection, College of Medical and Dental Sciences, University of Birmingham, B152IT, Edgbaston UK.

⁴German Center for Lung Research (DZL)


A



B



Supplementary Figure 1. CFSE-labeled microvesicles. MVs isolated from PLY (100 ng/ml, 4 hours)-treated A549 were labeled with CFSE. **a.** CFSE-MVs were analyzed by confocal microscopy. Brightfield (left panel), fluorescence (middle panel), and merged (right panel) of the CFSE-labeled MVs were taken. **b.** Representative FACS dot plot of CFSE-MVs stained with annexin V (APC) in the presence (Red dots) or absence of calcium (black dots in the lower left quadrant; negative staining).



Supplementary Figure 2. Full length blots.