

Supplementary information

Improvement in the long-term care burden after surgical treatment of patients with idiopathic normal pressure hydrocephalus: a supplementary study

Masatsune Ishikawa^{1,2*}, Shigeki Yamada^{3,2}, Masakazu Miyajima⁴, Hiroaki Kazui⁵, and Etsuro Mori⁶

¹Rakuwa Villa Ilios, Kyoto, Kyoto, Japan

²Normal pressure hydrocephalus Centre, Otowa Hospital, Kyoto, Kyoto, Japan

³Department of Neurosurgery, Shiga University of Medical Science, Otsu, Shiga, Japan

⁴Department of Neurosurgery, Juntendo Tokyo Koto Geriatric Medical Center, Kotoku, Tokyo, Japan

⁵Department of Neuropsychiatry, Kochi Medical School, Kochi University, Nankoku, Kochi, Japan

⁶Department of Behavioral Neurology and Neuropsychiatry, Osaka University United Graduate, Toyonaka, Osaka, Japan

Supplementary Table legends:

1. **Supplementary Table 1 [SuppleTable1]:** Independence level in disability in long-term care insurance system in Japan (LTCB disability)
2. **Supplementary Table 2 [SuppleTable2]:** Independence level in dementia in long-term care insurance system in Japan (LTCB dementia)
3. **Supplementary Table 3 [SuppleTable3]:** Model comparison between GLM and GLMM
4. **Supplementary data file 1 [SuppleData1]:** LTCB data in a wide format
5. **Supplementary data file 2 [SuppleData2]:** LTCB data in a long format