

**Temporal activity patterns suggesting niche partitioning of sympatric carnivores in Borneo,
Malaysia**

Miyabi Nakabayashi^{1*}, Tomoko Kanamori^{2,3}, Aoi Matsukawa⁴, Joseph Tangah⁵, Augustine
Tuuga⁶, Titol Peter Malim^{6†}, Henry Bernard⁷, Abdul Hamid Ahmad⁸, Ikki Matsuda^{4,7,9,10}, Goro
Hanya³

¹Graduate School of Advanced Science and Engineering, Hiroshima University, Hiroshima, Japan

²Japan Orangutan Research Center, Tokyo, Japan

³Primate Research Institute, Kyoto University, Aichi, Japan

⁴Wildlife Research Center, Kyoto University, Kyoto, Japan

⁵Forest Research Centre, Sabah Forestry Department, Sandakan, Sabah, Malaysia

⁶Sabah Wildlife Department, Kota Kinabalu, Sabah, Malaysia

[†]Titol Peter Malim passed away on 6th February 2021. His contributions as an author are listed in the Author Contributions section.

⁷Primate Studies-Borneo, Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Sabah, Malaysia

⁸Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Sabah, Malaysia

⁹Chubu University Academy of Emerging Sciences, Aichi, Japan


¹⁰Japan Monkey Centre, Aichi, Japan

*Corresponding author


E-mail: miyabi.nakabayashi@gmail.com

Phone number: +81-82-424-6930

Supplementary Figures



Supplementary Figure S1 Monthly numbers of cumulative photos of four carnivore species (banded civet, Malay civet, common palm civet and sun bear) and working cameras in the (a) DVCA, (b) LKWS, and (c) TWR. Dotted line and solid line indicate number of working cameras and cumulative photo number, respectively. Note that durations of more than three consecutive years are shown.



Supplementary Figure S2 Monthly numbers of cumulative species of carnivores in the (a) DVCA, (b) LKWS, and (c) TWR. Note that durations of more than three consecutive years are shown.