

Title:

Subchondral tibial bone texture of conventional X-Rays predicts total knee arthroplasty

Authors:

Ahmad ALMHDIE-IMJABBAR^{1,2}, Hechmi TOUMI^{1,2,3}, Khaled HARRAR⁴, Antonio PINTI^{1,5}, Eric LESPESSAILLES^{1,2,3}

Affiliations:

¹EA 4708- I3MTO Laboratory, University of Orleans, Orleans, France

²Translational Medicine Research Platform, PRIMMO, Regional Hospital of Orleans, France

³Department of Rheumatology, Regional Hospital of Orleans, France

⁴LIST Laboratory, University M'Hamed Bougara of Boumerdes, Algeria

⁵DeVisu-Design, Visuel, Urbain, EA 2445, UPHF, Valenciennes, France

E-mail addresses:

Ahmad ALMHDIE-IMJABBAR, (ahmad.almhdie@univ-orleans.fr), Hechmi TOUMI (hechmi.toumi@univ-orleans.fr), Khaled HARRAR (khaled.harrar@univ-boumerdes.dz), Antonio PINTI (antonio.pinti@uphf.fr), Eric LESPESSAILLES (eric.lespessailles@chr-orleans.fr)

Corresponding author:

Eric Lespessailles, Regional Hospital of Orleans, 14 avenue de l'Hôpital, 45067, Orleans Cedex 2, France

Supplementary Material




Figure S1: The most efficient TBT parameters chosen by the AIC algorithm for Scenario I. Parameters calculated in the micro- (left) and milli- (right) scales, in the horizontal (top) and vertical (bottom) directions




Figure S2: The most efficient TBT parameters chosen by the AIC algorithm for Scenario II. Parameters calculated in the micro- (left) and milli- (right) scales, in the horizontal (top) and vertical (bottom) directions