

Association between a dietary pattern high in saturated fatty acids, dietary energy density, and sodium with coronary heart disease

Nur Ain Fatinah Abu Bakar¹, Aryati Ahmad^{1,2*}, Wan Zulaika Wan Musa^{1&} Mohd Razif Shahril³, Nadiah Wan-Arfah¹, Hazreen Abdul Majid⁴, Carmen Piernas², Ahmad Wazi Ramli⁵, Nyi Nyi Naing⁶

Nutrients	Univariate analysis OR (95% CI)	p value
Energy intake (kcal/day)	1.001(1.001,1.002)	<0.001
Dietary energy density (DED) kcal/g	1.26(0.60, 2.65)	0.550
Fat % of EI	1.07(1.03, 1.11)	<0.001
SFA (g/day)	1.09(1.06, 1.11)	<0.001
SFA (g/1000 kcal)	1.10(1.04, 1.18)	0.003
Sodium (g/day)	1.00(1.00, 1.001)	<0.001
Sodium (mg/1000kcal)	1.00(1.00, 1.00)	0.553

Table S1. Simple logistic regression between single nutrients and CHD (n=365).

Binary logistic regression was used to test the association of each nutrient with CHD *Significant level at p-value <0.05. SFA: Saturated fatty acid. Abbreviations: CI: Confidence interval; DED: Dietary energy density; SFA: saturated fatty acids.