



# Supplemental figure 1



## Figure Legend: Supplemental figure 1

Symptom duration was compared between Asymptomatic/Mild group and Moderate/Severe group. The duration of the most prolonged symptoms from each participant was compared. There were no statistically significant differences by the Mann-Whitney U-test.

# Supplemental figure 2



## Figure Legend: Supplemental figure 2

Symptom duration was compared between ELISPOT-high group and low group. The duration of the most prolonged symptoms from each participant was compared. Although there were no statistically significant differences by the Mann-Whitney U-test, ELISPOT-low group showed a tendency for symptoms to persist for a longer period.

# Supplemental figure 3

|                                       | ELISPOT-high<br>(n=7) | ELISPOT-low<br>(n=8) |                |
|---------------------------------------|-----------------------|----------------------|----------------|
|                                       | Median (Q1-Q3)        |                      | <i>p</i> value |
| Sampling<br>(Days after<br>diagnosis) | 40 (35 - 46)          | 43 (38 - 117)        | <i>p</i> =0.61 |

## Figure Legend: Supplemental figure 3

The comparison of the number of days from diagnosis to the blood sampling between ELISPOT-high and -low group. The result showed no significant difference between those two groups with a p-value of 0.61 by the Mann-Whitney U test.