


Title: Summer diarrhea in children: a monocentric French epidemiological observational study.

Running title: Infectious summer diarrhea in children.


Authors: Camille Mallier, MD^{1,\$} & Elisa Creuzet^{2,\$}, Céline Lambert, MSc³, Julien Delmas, PharmD, PhD⁴, Audrey Mirand, PharmD, PhD^{5,6}, Emmanuelle Rochette, PhD^{1,7}, Stéphane Valot, PharmD^{8,9}, Maxime Moniot, PharmD², Frédéric Dalle, PharmD, PhD^{8,9,10}, Cécile Henquell, PharmD, PhD^{5,6}, Etienne Merlin, MD, PhD^{1,7}, Philippe Poirier, PharmD, PhD¹¹, Matthieu Verdan, MD^{1,\$} & Céline Nourrisson, PharmD, PhD^{11,\$,*}

Figure S1. Prevalence of each enteropathogen and co-detection according to the year.



EHEC: enterohemorrhagic *Escherichia coli*; EIEC: enteroinvasive *Escherichia coli*; EPEC: enteropathogen *Escherichia coli*; STEC: Shiga toxin-producing *Escherichia coli*.

Figure S2. Monthly distribution of the different infectious agents detected according to the year (a): bacteria (b), viruses (c) and parasites (d).



EHEC: enterohemorrhagic *Escherichia coli*; EIEC: enteroinvasive *Escherichia coli*; EPEC: enteropathogen *Escherichia coli*; STEC: Shiga toxin-producing *Escherichia coli*.