

Supplementary materials for:

***Meta*-Topolin–induced mass shoot multiplication and biosynthesis of valuable secondary metabolites in *Stevia rebaudiana* Bertoni bioreactor culture**

Agata Ptak^{1*}, Agnieszka Szewczyk², Magdalena Simlat¹, Alicja Błażejczak¹ & Marzena Warchoń³

¹Department of Plant Breeding, Physiology and Seed Science, University of Agriculture in Krakow, Łobzowska 24, 31-140 Krakow, Poland

²Department of Pharmaceutical Botany, Faculty of Pharmacy, Jagiellonian University Medical College, Medyczna 9, 30-688 Krakow, Poland

³The Franciszek Górski Institute of Plant Physiology, Polish Academy of Sciences, Niezapominajek 21, 30-239 Krakow, Poland

*Correspondence: mfptak@cyf-kr.edu.pl, agata.ptak@urk.edu.pl (A. Ptak)




Figure S1. HPLC chromatogram (analysis of steviol glycosides) of methanolic extract of *Stevia rebaudiana* (sample: in vitro shoots from 5 μ M of *meta*-Topolin) 1. rebaudioside A, 2. stevioside.




Figure S2. HPLC chromatogram (analysis of phenolic compounds) of methanolic extract of *Stevia rebaudiana* (sample: in vitro shoots from 5 μ M of *meta*-Topolin); 1. neochlorogenic acid, 2. chlorogenic acid, 3. isochlorogenic acid A, 4. isoquercetin, 5. rosmarinic acid, 6. quercitrin.