

Research Paper

Title: Genome-wide transcriptional profiling and physiological investigation elucidating the molecular mechanism of multiple abiotic stress response in *Stevia rebaudiana* Bertonni

Poonam Pal^{1,2}, Mamta Masand^{1,2}, Shikha Sharma^{1,2}, Romit Seth¹, Gopal Singh^{1,2}, Sanatsujat Singh¹, Ashok Kumar Yadav¹, Ram Kumar Sharma^{1,2*}

¹CSIR-Institute of Himalayan Bioresource Technology, Palampur -176061, India

²Academy of Scientific and Innovative Research (AcSIR), Ghaziabad-201002, India

***Correspondence:**

Dr. Ram Kumar Sharma

Biotechnology Division

CSIR-Institute of Himalayan Bioresource Technology

Palampur (H.P)

India

Email: rksharma.ihbt@gmail.com

Table ST1: Summary of Illumina generated sample-wise read information for 24 samples(raw read and filtered read) and mapping % of filtered reads on reference genome in *S. rebaudiana*. (C_L: Control_leaf; DS_L: Drought stress_leaf; SS_L: Salt stress_leaf; WS_L: Waterlogging stress_leaf; C_R: Control root; DS_R: Drought stress leaf; SS_R: Salt stress_root; WS_R: Waterlogging stress_root)

Sample detail	Raw reads	Filtered reads	Mapping% on the reference genome
C_L1	157208824	126844632	84.22%
C_L2	235205490	187570824	87.50%
C_L3	233621792	162834858	87.34%
DS_L1	179252334	139101150	84.24%
DS_L2	157111262	109811772	87.56%
DS_L3	158359182	126442772	86.02%
SS_L1	164293032	140056234	88.61%
SS_L2	156372390	125201916	87.26%
SS_L3	189000936	150060418	87.85%
WS_L1	214250770	146670206	88.17%
WS_L2	199847576	159675414	89.53%
WS_L3	163318122	134747470	84.84
C_R1	143765744	107512090	68.85%
C_R2	149920782	92979948	75.00%
C_R3	133804352	96784574	85.11%
DS_R1	125536220	111337902	87.24%
DS_R2	171486248	118066914	80.74%
DS_R3	188934556	135208036	79.99%
SS_R1	140451492	104335172	84.62%
SS_R2	161338114	106689310	85.11%
SS_R3	154342830	111977028	82.16%
WS_R1	162528914	130873462	75.05%
WS_R2	182677418	148969522	82.80%
WS_R3	209893032	162239074	78.19%
Total	4132521412	3135990698	83.66%

Table ST2: Summary of *de novo* assembly transcriptome of *S. rebaudiana*.

TRINITY Assembly Statistics	
No. of assembled transcripts	318859
Total length of transcript(bp)	274829027
Average sequence length (bp)	861
N50 length(bp)	1396
GC content	41.50%

Table ST3: List of unique abiotic stress-specific GO term processes enriched under drought stress (DS), salinity stress (SS) and waterlogging (WS)

GO term	Description	Hits	p-value
GO terms under drought stress (DS) in leaf			
GO:0006749	Glutathione metabolic process	26	0.003647
GO:0009896	Positive regulation of catabolic process	19	0.018135
GO:0009687	Absciscic acid metabolic process	13	0.017426
GO:0004575	Sucrose alpha-glucosidase activity	7	0.002712
GO:0005987	Sucrose catabolic process	5	0.018763
GO:0071218	Cellular response to misfolded protein	11	0.04302
GO:0009963	Positive regulation of flavonoid biosynthetic process	4	0.041248
GO:0072665	Protein localization to vacuole	11	0.020034
GO:0043269	Regulation of ion transport	13	0.028553
GO:0015081	Sodium ion transmembrane transporter activity	4	0.013339
GO:0046345	Absciscic acid catabolic process	4	0.000821
GO:0046345	Absciscic acid catabolic process	4	0.000821
GO terms under drought stress (DS) in root			
GO:0004024	Alcohol dehydrogenase activity, zinc-dependent	5	0.008174
GO:0005372	Water transmembrane transporter activity	21	0.000866
GO:0015925	galactosidase activity	10	0.049689
GO:1905421	Regulation of plant organ morphogenesis	8	0.026039
GO:1902584	Positive regulation of response to water deprivation	3	0.014925
GO:0010133	Proline catabolic process to glutamate	3	0.009575
GO:0009739	Response to gibberellin	27	0.016447
GO:0004553	Hydrolase activity, hydrolyzing O-glycosyl compounds	79	0.005714
GO:0015145	Monosaccharide transmembrane transporter activity	4	0.048927
GO:0017014	Protein nitrosylation	8	0.00126
GO:0051762	Sesquiterpene biosynthetic process	4	0.000947

GO:0051762	Sesquiterpene biosynthetic process	4	0.000947
GO terms under salinity stress (SS) in leaf			
GO:0007165	Signal transduction	16	0.036167
GO:0022804	Active transmembrane transporter activity	88	0.041043
GO:0009755	Hormone-mediated signaling pathway	104	0.000793
GO:0008194	UDP-glycosyltransferase activity	65	0.008685
GO:0015298	Solute:cation antiporter activity	10	0.026393
GO:0015368	Calcium:cation antiporter activity	5	0.023648
GO:0004672	Protein kinase activity	163	0.016796
GO:0006817	Phosphate ion transport	5	0.023152
GO terms under salinity stress (SS) in root			
GO:0004326	Tetrahydrofolylpolyglutamate synthase activity	3	0.031081
GO:0006206	Pyrimidine nucleobase metabolic process	4	0.013465
GO:2000104	Negative regulation of DNA-dependent DNA replication	5	0.019263
GO:0005666	RNA polymerase III complex	6	0.018018
GO:0045595	regulation of cell differentiation	11	0.041278
GO:0042575	DNA polymerase complex	7	0.003759
GO:0034399	Nuclear periphery	4	0.001346
GO:0006801	Superoxide metabolic process	6	0.046332
GO terms under waterlogging stress (WS) in leaf			
GO:0046394	Carboxylic acid biosynthetic process	36	0.038108
GO:0016597	Amino acid binding	3	0.006633
GO:0005524	ATP binding	18	0.012478
GO:0140657	ATP-dependent activity	17	0.000446
GO:0043168	Anion binding	60	0.046657
GO:0140657	ATP-dependent activity	17	0.000446
GO:0016053	Organic acid biosynthetic process	37	0.049448
GO:0006544	Glycine metabolic process	3	0.006633
GO:0030554	Adenyl nucleotide binding	20	0.007652
GO terms under waterlogging stress (WS) in root			

GO:0051787	Misfolded protein binding	17	0.016414
GO:0034220	Ion transmembrane transport	6	0.026483
GO:0008308	Voltage-gated anion channel activity	8	0.00437
GO:0070469	Respirasome	75	0.001814
GO:0016614	Oxidoreductase activity, acting on CH-OH	106	0.008426
GO:0015986	ATP synthesis coupled proton transport	18	0.000492
GO:0071555	Cell wall organization	73	0.003633
GO:0016052	Carbohydrate catabolic process	20	0.027918
GO:0098655	Cation transmembrane transport	118	0.005872

Table ST4. Detail of primers pairs used for validation using RT-qPCR

Transcript ID	Primer Sequence
TRINITY_DN123583_c1_g1_i2Fw	GCTGTTGTGGTTTTTACAAGTA
TRINITY_DN123583_c1_g1_i2Rw	CTAATCTCCCGTTCGTAATAAG
TRINITY_DN22036_c0_g3_i1Fw	AGCTATTTTTCAAGCCGTTAT
TRINITY_DN22036_c0_g3_i1Rw	AGATATGTGTGTGCGGTATGT
TRINITY_DN27941_c0_g1_i1Fw	TATCCTCCTGTGAGCAAACCT
TRINITY_DN27941_c0_g1_i1Rw	ATGTCGTGGTAATCCAATGT
TRINITY_DN115524_c1_g1_i1Fw	GGCTTCTCAGGTTGTTTTAAT
TRINITY_DN115524_c1_g1_i1Rw	AAGTTCCTTAACCTTCAACTCA
TRINITY_DN38046_c2_g2_i1Fw	GGCTCTAAAACCTAGCAAGATA
TRINITY_DN38046_c2_g2_i1Rw	AGTAGAACACCTAAGGGACATG
TRINITY_DN120851_c8_g2_i2Fw	GTATGTTTCTAACCGAGGAAGA
TRINITY_DN120851_c8_g2_i2Rw	AGTTGCTCTTTTAACACCTTGT
TRINITY_DN136789_c0_g1_i1Fw	GGAATGTTGGGGAAGATCAA
TRINITY_DN136789_c0_g1_i1Rw	ACCAATGGGAGATGGTTGAC
TRINITY_DN104322_c0_g2_i2Fw	GATATGTGAGTGATGTTTGGA
TRINITY_DN104322_c0_g2_i2Rw	TGAGGGATTACCTTATCCTTA
TRINITY_DN25856_c0_g1_i2Fw	CAATGGACAGATCAAATTACAA
TRINITY_DN25856_c0_g1_i2Rw	CATCCTCCATAATCGTCTTAAT
TRINITY_DN118955_c1_g1_i3Fw	ACTGGGTAAACGGGCCAATA
TRINITY_DN118955_c1_g1_i3Rw	ACGAGCAAAGTGTTTGAAGAGA
TRINITY_DN100891_c0_g1_i1Fw	CGGATTAAAAGCTAGAGAAAGA
TRINITY_DN100891_c0_g1_i1Rw	CTTGAATATGATCTGGGATTTT
TRINITY_DN113546_c0_g1_i1Fw	CTCAAGGAATTGTACCTTTTGT
TRINITY_DN113546_c0_g1_i1Rw	CACAAACCAAGAATGATGTAGT
TRINITY_DN114691_c0_g2_i1Fw	TGCTGAGAAAACACACATACA
TRINITY_DN114691_c0_g2_i1Rw	CGTCACTTGGGGTAACTG

Figure S1. Functional annotation of the *de novo* assembled transcripts (a) and KEGG enrichment of differentially expressed transcripts (DEGs) in leaf and root tissues (b) under abiotic stress.








Figure S2. Gene ontology (GO) enrichment analysis. Dot plot shows GO terms upregulated and downregulated identified under **(a)** drought (DS_L and DS_R), **(b)** salinity (SS_L and SS_R), and waterlogging (WS_L and WS_R) stress using g:Profiler to be enriched (adjusted p-value < 0.05) among the genes. The size of the dots is based number of genes enriched in the pathway, and the color of the dots represents the adjusted p-values





(b)





(c)




Figure S3. Differentially expressed transcriptional factor families under DS, SS and WS in *S. rebaudiana*. (a) Top 10 TF families expressed under DS, SS and WS. (b) The number of transcripts (up and downregulated) and respective Venn diagram representing shared (upregulated) TF among stresses in leaf and root tissues. (c) Heatmap analysis of differentially expressed shared TF encoding genes under DS, SS and WS.




Figure S4. Heatmap analysis of differential upregulated transcripts encoding for TF unique to each abiotic stress in *S. rebaudiana*.




Figure S5. The qRT-expression analysis of the selected key significant DEGs in abiotic stress treated multiple genotypes. The columns indicate the qRT-PCR value under DS (Red) and SS (green), in three genotypes. Each bar represents the mean $\pm$ SD of triplicate assays.

