

Comparison of three primer pairs for molecular sex determination in Eurasian pygmy owls (*Glaucidium passerinum*)

Simona Stehlíková Sovadinová^{1*}, Chahrazed Mekadim², Erkki Korpimäki³, Jakub Mrázek², Marek Kouba^{1,4}

¹Czech University of Life Sciences Prague, Faculty of Agrobiological Sciences, Department of Ethology and Companion Animal Science, Kamýcká 129, Praha 6 – Suchbátka, 165 00, Czechia

²Czech Academy of Sciences, Institute of Animal Physiology and Genetics, Laboratory of Anaerobic Microbiology, Václavská 1083, Praha 4 – Krč, 142 20, Czechia


³University of Turku, Department of Biology, Section of Ecology, 20014 Turku, Finland

⁴Czech University of Life Sciences Prague, Faculty of Forestry and Wood Sciences, Department of Game Management and Wildlife Biology, Kamýcká 129, Praha 6 – Suchbátka, 165 00, Czechia


*Corresponding author: s.stehlikova.s@gmail.com

ORCID – Simona Stehlíková Sovadinová: 0009-0008-9431-5156; Chahrazed Mekadim: 0000-0002-5793-3501; Erkki Korpimäki: 0000-0001-7596-1955; Jakub Mrázek: 0000-0003-1907-3333; Marek Kouba: 0000-0003-2262-5733


Figures




Supplementary Fig. 1. Original agarose gel from electrophoresis of PCR products of Eurasian pygmy owls using primer set CHD1F/CHD1R.



Supplementary Fig. 2. Melt curve analysis of quantitative polymerase chain reaction products from Eurasian pygmy owls using primer set P2/P8. The curve was identical for males and females.



Supplementary Fig. 3. Original agarose gel from electrophoresis of PCR products of Eurasian pygmy owls using primer sets P2/P8 (upper) and 2550F/2718R (lower).



Supplementary Fig. 4. Melt curve analysis of quantitative polymerase chain reaction products from Eurasian pygmy owls using primer set 2550F/2718R. The curve was identical for males and females.