




*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.




*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.




*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.




*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.




*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.




*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.




*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.



*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.



*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.



*The blots were cut prior to hybridization with antibodies during blotting. Some membranes were washed and reprobed to detect the unphosphorylated proteins or other proteins with similar molecular weight. The overexposed parts (red color) are due to the reprobe reasons and were not used in data analysis.

