

WWP1 localizes in the Golgi apparatus and contributes to maintaining glycosaminoglycan synthesis in adipocytes

Yuka Nozaki^{1,6}, Fumika Suwa^{1,6}, Kazuhiro Furuya¹, Masahiro Komeno¹, Shunsuke Hoshino², Yuhei Mizunoe¹, Kyohei Higashi¹, Masaki Kobayashi^{3,4*}, Yoshikazu Higami^{1,5*}

1. Faculty of Pharmaceutical Science, Tokyo University of Science, Chiba 278-8510, Japan

2. Molecular Glycobiology, Research Team for Mechanism of Aging, Tokyo Metropolitan Institute for Geriatrics and Gerontology, Tokyo 173-0015, Japan

3. Department of Nutrition and Food Science, Graduate School of Humanities and Sciences, Ochanomizu University, Tokyo 112-8610, Japan


4. Institute for Human Life Innovation, Ochanomizu University, Tokyo 112-8610, Japan

5. Division of Cell Fate Regulation, Research Institute for Biomedical Science, Tokyo University of Science, Chiba 278-8501, Japan

6. The authors contributed equally


*Correspondence: higami@rs.tus.ac.jp, kobayashi.masaki2@ocha.ac.jp

Supplemental Figure 1



Supplemental Figure 1. WWP1 influenced endoplasmic reticulum stress in differentiated 3T3L1 adipocytes. (A-D) The mRNA level of *Grp78*, *Atf4*, *Total Xbp1* and *Spliced Xbp1* in differentiated mock and WWP1 overexpressing (OE) differentiated 3T3L1 adipocytes were analyzed by real-time PCR. *Rps18* was used as a reference control. Differences between each value were analyzed by Student's t-test. (*: $p < 0.05$, ***: $p < 0.001$).


Supplemental Figure 2




Supplemental Figure 2. Expression level of endogenous WWP1 in differentiated 3T3L1 adipocytes under Golgi stress. (A) The differentiated 3T3L1 adipocytes were treated by DMSO or 10 μ M monensin for 6 hours and analyzed by immunoblotting for anti-WWP1 antibodies. Lamin B1 was used as a loading control. (B) The quantitative data are shown as the mean $\pm$ SD (n = 3). Differences between each value were analyzed by Student's t-test.

Supplemental Figure 3

Unsaturated disaccharide unit of heparan sulfate



Unsaturated disaccharide unit of chondroitin sulfate



Supplemental Figure 3. The structure of unsaturated disaccharides of HS and CS.