

Supplementary Table 1. Examination performed prior to FDG-PET/CT

Examination	Total	FUO (n = 21)	IUO (n = 24)	p-value
<i>Laboratory test</i>				
CBC	45 (100%)	21 (100%)	24 (100%)	N/A
Hepatic function	45 (100%)	21 (100%)	24 (100%)	N/A
Renal function	45 (100%)	21 (100%)	24 (100%)	N/A
Electrolytes	45 (100%)	21 (100%)	24 (100%)	N/A
Viral tests	19 (42%)	13 (62%)	6 (25%)	0.012
ANCA	34 (76%)	17 (81%)	17 (71%)	0.43
ANA	37 (82%)	18 (86%)	19 (79%)	0.57
Urinalysis	41 (91%)	21 (100%)	20 (83%)	0.050
Blood culture	34 (76%)	16 (76%)	18 (75%)	0.93
Peripheral blood smear	2 (4%)	0 (0%)	2 (8%)	0.18
Bone marrow exam	5 (11%)	4 (19%)	1 (4%)	0.11
<i>Imaging test</i>				
Chest radiography	28 (62%)	11 (52%)	17 (71%)	0.20
Abdominal US	9 (20%)	3 (14%)	6 (25%)	0.37
Echocardiography	3 (7%)	1 (5%)	2 (8%)	0.63
EGD	8 (18%)	5 (24%)	3 (12%)	0.32
CS	6 (13%)	2 (10%)	4 (17%)	0.48
CT	27 (60%)	12 (57%)	15 (62%)	0.71
MRI	5 (11%)	1 (5%)	4 (17%)	0.20

CBC: complete blood count; ANCA: antineutrophilic cytoplasmic antibodies; ANA: antinuclear antibody; US: ultrasonography; TTE: trans-thoracic echocardiology; EGD: esophagogastroduodenoscopy; CS: colonoscopy; CT: computed tomography; MRI: magnetic resonance imaging. All variables were compared using Pearson's chi-square test/Fisher's exact test.