

Vulture dominance in a scavenger assemblage in the neotropical dry forest

Paul Monar-Barragán^{1,2*}, Félix Hernán Vargas², Rubén Pineida¹, Andy Salazar¹, Emilio Virgos³, Fabricio Narváez¹, Sebastián Kohn¹, Juan de Dios Morales⁴, Juan Sebastián Restrepo-Cardona^{1,5}

¹Fundación Cóndor Andino Ecuador, Quito-Pichincha, Ecuador. ²The Peregrine Fund-Neotropical Program, Santa Cruz-Galápagos, Ecuador. ³Rey Juan Carlos University, Móstoles-Madrid, Spain. ⁴The Wild GYE Initiative, Guayaquil-Guayas, Ecuador. ⁵Department of Wildlife Ecology and Conservation, University of Florida, Gainesville-FL, USA.

*Email: pmonar@fundacioncondor.org

Supplementary Table 2. Generalized linear mixed models (GLMM) of mammalian facultative scavenger community variables. Significant P-value (<0.05) highlighted in bold.

Response variable	Model	Estimate	std error	p-value
Mammal species richness	(Intercept)	-0.4030	0.37260	0.2794
	Carcass size	-0.6150	0.41180	0.1353
	Season	-0.8830	0.47470	0.0629
Mammal abundance	(Intercept)	-0.5826	0.46860	0.2138
	Carcass size	-0.6119	0.36140	0.0904
	Season	-0.7690	0.61120	0.2083
<i>Eira barbara</i> abundance	(Intercept)	-13.465	0.72090	0.0618
	Carcass size	-12.272	0.80850	0.1290
	Season	-13.713	0.89850	0.1269
<i>Leopardus pardalis</i> abundance	(Intercept)	-14.031	0.52970	0.0080
	Carcass size	-0.4633	0.43720	0.2892
	Season	-0.1333	0.61760	0.8290
<i>Canis lupus familiaris</i> abundance	(Intercept)	-88.840	7,502	0.2360
	Carcass size	<0.0001	1,017	1
	Season	-620.50	>100000	1