

**Changes in blood concentration of tranexamic acid in dialysis and non-dialysis patients undergoing cardiac surgery: a prospective observational cohort study**

Takahiro Tamura<sup>1</sup>, Tatsuro Yokoyama<sup>2</sup>, Tsuyoshi Nakai<sup>3</sup>, Yasuhiro Miyagawa<sup>3</sup>, Kimitoshi Nishiwaki<sup>1</sup>

<sup>1</sup> Department of Anesthesiology, Nagoya University Graduate School of Medicine, Nagoya 4660550, Japan

<sup>2</sup> Department of Perioperative Management System, Nagoya University Graduate School of Medicine, Nagoya 4660550, Japan

<sup>3</sup> Department of Pharmacotherapeutics and Informatics, Fujita Health University School of Medicine, Toyoake, Aichi 4701192, Japan

**Supplementary Table S1.** Classification of chronic kidney disease in Japanese patients  
by estimated glomerular filtration rate (eGFR)

| CKD stage | eGFR (mL/min/1.73 m <sup>2</sup> ) |
|-----------|------------------------------------|
| G1        | ≥ 90                               |
| G2        | 60–89                              |
| G3a       | 45–59                              |
| G3b       | 30–44                              |
| G4        | 15–29                              |
| G5        | < 15                               |

eGFR: estimated glomerular filtration rate; CKD: chronic kidney disease