

**Molecular insights on the proangiogenic effects of VEGF like growth factor derived from
horseshoe crab perivitelline fluid**

K. G. Aghila Rani ^{1, #}, Hrishikesh Pandit ^{1, #}, Rambhadur Subedi ^{1, #}, Sayli Otiv-Pandit ¹, Anil Bankati Jogdand ¹, Hajra Gupta ¹, Prayagraj Fandilolu ², Sushama Rokade ¹, Susan Idicula-Thomas ², Anil Chatterji ³, Taruna Madan ^{1, 4, *}

¹ Department of Innate Immunity, National Institute for Research in Reproductive and Child Health (ICMR), J. M. Street, Parel, Mumbai 400 012, India.


² Biomedical Informatics Center, ICMR-National Institute for Research in Reproductive and Child Health, J. M. Street, Parel, Mumbai 400 012, India.

³ Malkolak Institute of Marine Sciences, 8 Astha Society, Bella Vista Colony, Dona Paula, Goa 403 004, India.

⁴ ICMR-Division of Development Research (Govt of India), New Delhi, 110029, India.

Supplementary Figure 1:

Full length silver-stained SDS-PAGE gel (raw data of Fig. 1A) showing crude PVF (L3) along with protein ladder (L4). L1 and L2 show profile of other samples ran alongside crude PVF.



Supplementary Figure 2:

Full length silver-stained SDS-PAGE gel (raw data of Fig. 1C) revealing profile of PVF fractions. L1 (protein ladder), L2 (fraction II), L3 (fraction III), L4 (fraction IV), L5 (fraction V), L6 (fraction VI), L7 (fraction VII), L8 (fraction VIII), L9 (protein ladder).

