

Effects of soil biofumigation on non-target springtails (Collembola) and earthworms
(Opisthopora)

Usha Panta^{1*}, Ashley Jernigan², and Arash Rashed^{1*}

¹ Department of Entomology, Southern Piedmont Agricultural Research and Extension Center,
Virginia Tech, Blackstone, VA, USA

² School of Plant and Environmental Sciences, Virginia Tech, Blacksburg, VA, USA

*Corresponding authors: Usha Panta, usha@vt.edu;

Arash Rashed, arashr@vt.edu

Supplementary Materials

Fig. S1: Imaging setup used for the springail number quantification using a Canon EOS Rebel T7 camera equipped with an EF-S 18-55 lens mounted on a tripod.




Fig. S2: Two-way Polyvinyl Chloride (PVC) olfactometer with central pipe (A), side pipe (B), and elbow piece (C).



Fig. S3: Cloth mesh pouches filled with sterilized potting mix and springtails.



Fig. S4: Mean ($\pm$ SE) number of springtails in control and brown mustard seed meal (BMSM)-treated pots using untransformed data across all time points (1 to 40). The main graph presents the complete time course, while the inset highlights days 1 to 19 to emphasize initial treatment effects. Asterisks indicate significant differences ($p < 0.05$) between treatments at each time point.

