

Supplementary Table 2: Age, sex, *T. gondii* infection and inflammatory marker concentrations between cases and healthy controls

Characteristic	Overall N=175	Cases; n=160	Healthy controls N=15	p-value
Age				0.3
<i>Mean (SD)</i>	34(13)	34(14)	30(8)	
<i>Median (Min, Max)</i>	31 (8, 75)	32(8,75)	29(22, 56)	
Sex				0.2
<i>Male</i>	19 (11%)	16 (10%)	3(20%)	
<i>Female</i>	156 (89%)	144 (90%)	12(80%)	
<i>T. gondii</i>				0.006
<i>Mean (SD)</i>	0.45(0.7)	0.5(0.72)	0.0(0.0)	
<i>Median (Min, Max)</i>	0.0(0.0, 1.93)	0.0(0.0, 1.93)	0.0(0.0)	
IL-10 (pg/mL)				<0.001
<i>Mean (SD)</i>	236(65)	255(22)	37(5)	
<i>Median (Min, Max)</i>	249(32, 320)	254(222, 320)	37(32, 45)	
IL-1β (pg/mL)				<0.001
<i>Mean (SD)</i>	28(9)	30(7)	7(1)	
<i>Median (Min, Max)</i>	28(6, 49)	29(21, 49)	7(6, 10)	
IL-6 (pg/mL)				<0.001
<i>Mean (SD)</i>	173(120)	189(114)	10(1)	
<i>Median (Min, Max)</i>	147(9, 762)	149(107, 762)	10(9, 12)	
INF-γ (pg/mL)				<0.001
<i>Mean (SD)</i>	81(18)	86(7)	28(5)	
<i>Median (Min, Max)</i>	83(22, 100)	85(64, 100)	28(22, 43)	
TNF-α (pg/mL)				<0.001
<i>Mean (SD)</i>	71(26)	77(17)	6(1)	
<i>Median (Min, Max)</i>	75(5, 100)	77(24, 100)	6(5, 7)	