

Southward decrease in the protection of persistent giant kelp forests in the northeast Pacific

Authors: Nur Arafeh-Dalmau^{1,2*}, Kyle C. Cavanaugh³, Hugh P. Possingham¹, Adrian Munguia-Vega^{4,5}, Gabriela Montaña-Moctezuma⁶, Tom W. Bell⁷, Kate Cavanaugh³, Fiorenza Micheli⁸.

Affiliations:

¹Centre for Biodiversity and Conservation Science, School of Biological Sciences, The University of Queensland, St Lucia, Queensland, Australia.

²School of Earth and Environmental Sciences, The University of Queensland, St Lucia, Queensland, Australia.

³Department of Geography, University of California Los Angeles, Los Angeles, CA, USA.

⁴Desert Laboratory on Tumamoc Hill, University of Arizona, Tucson, AZ, 85721, USA.


⁵@Lab Applied Genomics, La Paz, Baja California Sur, Mexico.

⁶Instituto de Investigaciones Oceanológicas, Universidad Autónoma de Baja California, Ensenada, Mexico.


⁷Department of Applied Ocean Physics and Engineering, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, 02543

⁸Hopkins Marine Station and Stanford Center for Ocean Solutions, Pacific Grove, CA 93950, USA.

*Correspondence to: nuracons@gmail.com



Supplementary Figure 1: The spatial distribution of kelp by level of persistence and marine protected areas by level of protection in the Northeast Pacific Ocean. We provide fine-scale examples for each region.



Supplementary Figure 2: Boxplot with the occupancy value of detected kelp pixels ($n = 408,906$) for each of the four regions in the Northeast Pacific Ocean.