

Supplementary Table 1

Zero-order correlations between demographics, childhood maltreatment, and clinical symptoms

Variable	1	2	3	4	5	6	7	8	9	10
1. Age	1									
2. Gender* (female)	.27	1								
3. Physical abuse	.28	.01	1							
4. Emotional abuse	.27	.07	.23***	1						
5. Sexual abuse	.29	.30***	.05	-.03	1					
6. Physical neglect	.34	.03	.17***	.11**	.19***	1				
7. Emotional neglect	.33	.10**	.11***	.25***	.08*	.18***	1			
8. PHQ-9	-.18***	.05	.19	.23	.27**	.26*	.28**	1		
9. PC-PTSD	-.15***	.01	.15*	.15**	.19***	.22***	.23**	.57***	1	
10. GAD-7	-.24***	.11**	.22	.26**	.21	.23	.27**	.83***	.56***	1

Note. Gender: 0 = male, 1 = female; PHQ-9 = depressive symptoms; PC-PTSD = PTSD symptoms; GAD-7 =general anxiety

* $p < .05$; ** $p \leq .01$; *** $p \leq 0.001$

Supplementary Table 2

Model fit indices results for the SEM analyses comparing the two models, controlled for confounding variables

Model	Symptoms	RMSEA	SRMR	CFI	TLI	AIC	BIC
Abuse/Neglect Approach	PTSD	0.042	0.047	0.819	0.739	15742	15839
	Depression	0.042	0.047	0.827	0.751	17379	17467
	Anxiety	0.042	0.047	0.847	0.78	17188	17275
Dimensional CM Approach	PTSD	0.053	0.05	0.624	0.454	14268	14373
	Depression	0.053	0.05	0.633	0.467	15909	16014
	Anxiety	0.053	0.05	0.662	0.509	15725	15830

SRMR: Standardized Root Mean Square Residual

CFI: comparative fit index

RMSEA: Root Mean Square Error of Approximation

Supplementary Table 3

Model Fit Indices and R-squared Results for OLS Regression Analyses

Top Predictors	Symptoms	R-squared	AIC	BIC
Emotional Neglect, Sexual Abuse	PTSD	0.1485	2390	2414
Physical Neglect, Emotional Abuse	Depression	0.1056	3991	4114
Emotional Abuse, Sexual Abuse	Anxiety	0.949	3828	2951

Note. AIC = Akaike Information Criterion; BIC = Bayesian Information Criterion

Predictors inserted in the models were 'Physical Abuse', 'Emotional Abuse', 'Sexual Abuse', 'Emotional Neglect' and 'Physical Neglect'