

Protein serine/threonine phosphatase PPEF-1 suppresses genotoxic stress response via dephosphorylation of PDCD5

Soo-Yeon Park^{1,7}, Jaesung Seo^{1,7}, Hyo-Kyoung Choi^{2,7}, Hye-Jeong Oh¹, Garam Guk¹, Yoo-Hyun Lee³, Jeongmin Lee⁴, Woo Jin Jun⁵, Kyung-Chul Choi^{6,*} and Ho-Geun Yoon^{1,*}

¹Department of Biochemistry and Molecular Biology, Brain Korea 21 PLUS Project for Medical Sciences, Yonsei University College of Medicine, Seoul, South Korea

²Division of Nutrition and Metabolism Research Group, Korea Food Research Institute, Gyeonggi-do, South Korea

³Department of Food and Nutrition, The University of Suwon, Kyunggi-do 445-743, South Korea

⁴Department of Medical Nutrition, Kyung Hee University, Kyunggi-do, 446-701, South Korea

⁵Department of Food and Nutrition, Chonnam National University, Gwangju, South Korea

⁶Department of Biomedical Sciences, and Department of Pharmacology, University of Ulsan College of Medicine, Seoul, South Korea

⁷These authors contributed equally to this work.

*Corresponding author. *E-mail:* choikc75@amc.seoul.kr (KCC) or yhgeun@yuhs.ac (HGY)

Fig. 1a








Fig. 1b




Fig. 1c




Fig. 1d




Fig. 2a




Fig. 2b




Fig. 2c




Fig. 2e




Fig. 2f





Fig. 2g




Fig. 3b




Fig. 3c




Fig. 4a




Fig. 4c




Fig. 4d




Fig. 4e




Fig. 5c

