

**Agreement of patient-measured intraocular pressure using rebound tonometry
with Goldmann applanation tonometry (GAT) in glaucoma patients**

^{1,6}Shaoying TAN, Ph.D., MD,

^{1,4}Marco YU, PhD,

^{1,2}Nafees BAIG, MRCS, FCOphth(HK),

^{1,2,5}Linda HANSAPINYO, MD,

^{1,2,3}Clement C THAM, FRCS, FCOphth(HK)

¹Department of Ophthalmology and Visual Sciences, The Chinese University of Hong Kong, Hong Kong;

²Hong Kong Eye Hospital, Kowloon, Hong Kong;

³Department of Ophthalmology and Visual Sciences, Prince of Wales Hospital, Shatin, Hong Kong.

⁴Department of Mathematics and Statistics, Hang Seng Management College, Hong Kong;

⁵Department of Ophthalmology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand;

⁶Department of Ophthalmology, Chinese PLA General Hospital, Beijing, China.

Patient Questionnaire on safety and ease of use of ICare rebound self-tonometry

Do you find instructions of the manual clear?

☐ Very unclear ☐ Unclear ☐ Fair ☐ Clear ☐ Very Clear

Do you find instructions of the manual adequate?

☐ Very Inadequate ☐ Inadequate ☐ Fair ☐ Adequate ☐ Very Adequate

Do you find the ICare rebound self-tonometry easy to use?

☐ Very Difficult ☐ Difficult ☐ Fair ☐ Easy ☐ Very Easy

Do you find the ICare rebound self-tonometry safe to use?

☐ Very Unsafe ☐ Unsafe ☐ Fair ☐ Safe ☐ Very Safe

Have you ever had any eye injury while using ICare rebound self-tonometry?

☐ Yes ☐ No

If yes, please write down the detail of injury.
