

The impact of the RBM4-initiated splicing cascade on modulating the carcinogenic signature of colorectal cancer cells

Jung-Chun Lin¹, Yuan-Chii Gladys Lee^{2,*}, Yu-Chih Liang^{1,*}, Yang C. Fann³, Kory R. Johnson³, Ying-Ju Lin⁴

¹School of Medical Laboratory Science and Biotechnology, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan

²Graduate Institute of Biomedical Informatics, Taipei Medical University, Taipei, Taiwan

³Information Technology and Bioinformatics Program, Division of Intramural Research, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA

⁴School of Chinese Medicine, China Medical University, Taichung, Taiwan

*These authors contributed equally to this work.

Running Title: RBM4a modulates CRC-related splicing events

To whom correspondence should be addressed: Jung-Chun Lin

School of Medical Laboratory Science and Biotechnology, College of Medical Science and Technology, Taipei Medical University

250 Wu-Hsing Street, Taipei 11031, Taiwan

Telephone: +886-2-27361661 ext. 3330

Fax: +886-2-27324510

E-mail: lin2511@tmu.edu.tw

Supplemental Table 1. PCR Primers.

Gene	Forward	Reverse
Nova1-AS	tgccccaaatgtggccaagaca	agccttcacagtagcacctccct
Nova1-Total	tctgatcccatgaccacctcca	ctgctggaaaggccgcaaca
SRSF6-AS	acgacgccgtttacga	ttaaatcttgccaactgcac
SRSF6 mini (F2)	atccagctgagctttatgctgtattg	atcctgcagtaagggaagtcaaataagac
Gapdh	cggagtcaacggatttggtcgtatg	agccttctccatggtggtgaagac
VEGF165-AS	ttgtacaagatccgcagacg	gttctgtatcagtctttcctgg
SV40	tttggaggcctaggctttt	
E-cadherin	tggaggaattcttgctttgc	cgtacatgtcagccagcttc
N-cadherin	ggcttaatggtgattttgctc	ctcaagtcatagtcctggtc

Supplemental Table 2. Quantitative RT-PCR Primers.

Gene	Forward	Reverse
Nova 1	tgagggaacaggtggagaac	agctggacacggagcttta
E-cadherin	cagaaagtttccacaaa	aaatgtgagcaattctgctt
N-cadherin	cgagccgcctgcgctgccac	cgtgtctctccgtccccgc
Gapdh	aaggtcatcccagagtgaa	ctgcttcaccaccttcttga