

Table S4. Sequences of primers used for semi-quantitative RT-PCR and qPCR.

Gene	Forward Primer (5' to 3')	Reverse Primer (5' to 3')
Semi-quantitative RT-PCR primers		
<i>BghEfc1</i>	CCAACTCGAAGTATAGATCCAAAAT	GTGAACTTCCTTCCGCTGG
<i>BghEfc2</i>	ACACCCATAAATTC AATTCGT	AAGCTCAAACAAGATGGA ACTA
<i>BghEfc3</i>	CACTCCAAACGCAAACATCACA	TAGCCTGGGTGTGCGAGG
<i>BghEfc4</i>	AGCTGCTACTTCCTATCTGCA	TTAGATACACTCTTCGGGCCGA
<i>BghEfc5</i>	TCCGTGAGACGAAATTCTCG	GGAAGGGACATAGTCGCCA
<i>BghEfc6</i>	CCACACATTTTAAATAGTCGACC	AGCCATTTACAACGTCGCATG
<i>BghEfc7</i>	CGCGTCGTTATTCAGCTGTC	CATCTTTATTCTTGTTGCGCCTT
<i>BghEfc8</i>	GCCTATTTTCCTTGGCCCTA	CTCGGCGACTTCCTCTTCA
<i>BghEfc9</i>	AAGTGGTTCATAGCGTTCATGT	TTCCTAAACGCGATCTGTCTAA
<i>BghEfc10</i>	GTTGAGGCTGGATATTTGTGTC	CCTATAACCCCTGCCGCT
<i>BghEfc12</i>	CACCATGTATTGGGATTGTGATGATTATAG	CCAGCCTAGTCTCTGAAC
<i>BghEfc15</i>	CACCATGTTGGATGGTTACCGATGTG	TGCTATGCATGCCATTAGC
<i>BghEfc16</i>	CACCATGTCATACTATGACTGCAACGG	CTCAATACATATATTTGCATTCTG
<i>BghEfc18</i>	CACCATGATGATAGGCTATCAATGTGAC	CCGTATCATTAAATTAATGGCAC
<i>BghEfc19</i>	CACCATGGAACAATATTTTAAATGTCATTC	GTTAATTCCGGTGGAAG
<i>BghEfc21</i>	CACCATGGCGGATTATCGTTGTGATAAT	CAGGTTTATTGACAGCTAATTAAG
<i>BghEfc26</i>	CACCATGCGAACGTGGCAATGCCG	CTGCCTAGGTGCATTCTTC
<i>BghEfc27</i>	ATCATTAAAGTACGGCTGACGA	AATGTCCTTACACGGTTCGT
<i>BghEfc29</i>	CACCATGGTGCAGTGTGTTTGGTGAC	GCCGACTAATACGGGC
<i>BghEfc30</i>	CACCATGTTAATGACTTATGAATGTGTGAG	GTGTTATGAGGCCAGCC
<i>BghEfc35</i>	CACCATGGCCTTCAAGGGGTATCATA	TGTTACCGAACAATGCCAC
<i>BghEfc36</i>	CACCATGCTGGATGGTTACAATTGTCT	CGCATATCATTCAATAATTTT
<i>BghEfc37</i>	GCGCCTGGAGAGTATGTTTA	TAAGGATAGGGCACGGCA
<i>BghHistone H3</i>	CGACTTGCGGTTCCAGTCA	ACACAAATTGGTATCCTCGAAGAGT
<i>Bghβ-tubulin</i>	CAGCAGATGTTTGATCCGAA	AGAACATAGGGCAGTTTGA
qPCR primers		
<i>BghEfc1</i>	GCTTTTCTTAAGCCATTTGA	TTCGTTTCAACCTCTTCTTC
<i>BghEfc2</i>	CATCTTGTTTGAGCTTGTC A	CCATACTACCCCAAATAGCA
<i>BghEfc3</i>	ACAGAGCTGTCGCTAGATT C	TTTCTATTGGCTGTTCTTA
<i>Bghβ-tubulin</i>	ATCGTATCCGCTACTTCAGA	CTGAGGCCGAGTCTAATATG
<i>HvGAPDH</i>	GTGAGGCTGGTGCTGATTACG	TGGTGCAGCTAGCATTGAGAC