

<i>Clostridium</i> sp.	F ^a	Restriction Enzyme (HaeIII) digestion fragments (nucleotides, nts)																	
<i>C. chauvoei</i>	6/8	■	78		■	22	■						■						
<i>C. colicanis</i>	9/9					■	22	■	573			■	312	■	144	■			
<i>C. botulinum</i>	32/128	■	78		■	22	■	572			■	456			■				
<i>C. butyricum</i>	9/32					■	22	■	85	■	487		■	457			■		
<i>C. butyricum</i>	21/32					■	22	■	572			■	456			■			
<i>C. acetobutylicum</i>	6/24					■	22	■	572			■	454			■			
<i>C. beijerinckii</i>	18/23					■	22	■	572			■	454			■			
<i>C. baratii</i>	7/8					■	22	■	571			■	456			■			
<i>C. sardiniense</i>	9/9					■	22	■	571			■	456			■			
<i>C. botulinum</i>	79/128					■	595			■	455			■	77		■		
<i>C. sporogenes</i>	7/11					■	595			■	455			■	77		■		
<i>C. tetani</i>	8/8	■	78		■	594			■	456			■	77		■	■		
<i>C. botulinum</i>	4/128					■	595			■	455			■					
<i>C. pasteurianum</i>	2/13					■	595			■	455			■					
<i>C. pasteurianum</i>	8/13					■	595			■	455			■	71	■	7	■	
<i>C. pasteurianum</i>	2/13					■	595			■	105	■	350	■					
<i>C. subterminale</i>	2/8			■	44	■	107		■	488	■	454			■	77		■	
<i>C. subterminale</i>	4/8					■	107		■	488	■	456			■	77		■	
<i>C. kluyveri</i>	14/14	■	78		■	1126										■			
<i>C. perfringens</i>	63/92			■	44	■	22	■	1104										■
<i>C. perfringens</i>	24/92			■	44	■	22	■											
<i>C. acetobutylicum</i>	16/24	■	399																■
<i>C. botulinum</i>	8/128	■ NA																	
<i>C. novyi</i>	17/17	■ NA																	

^aFrequency of organisms showing this R.E. digestion pattern.

Symbol (Filled square) indicates RE site in the 16S rDNA sequences. NA indicates, not applicable.