

Family Medicine Clinics Affiliated with St. Michael's Hospital, the University of Toronto (n=4)

Clinics were matched (1 large, 1 small) and randomized

Intervention Group

Control Group

Clinic 1

Clinic 2

Clinic 3

Clinic 4

Pre-Intervention

Pre-Intervention

Pre-Intervention

Pre-Intervention

Total Sample (n=2706)
Allocated to study* (n=119)

Total Sample (n=6571)
Allocated to study* (n=209)

Total Sample (n=1298)
Allocated to study* (n=122)

Total Sample (n=3607)
Allocated to study* (n=247)

Analyzed (n=95)
Excluded from analysis:
Chart not found (n=14)
Not meeting inclusion
criteria (n=10)

Analyzed (n=153)
Excluded from analysis:
Chart not found (n=23)
Not meeting inclusion
criteria (n=33)

Analyzed (n=94)
Excluded from analysis:
Chart not found (n=18)
Not meeting inclusion
criteria (n=10)

Analyzed (n=167)
Excluded from analysis:
Chart not found (n=19)
Not meeting inclusion
criteria (n=61)

Post-Intervention

Post-Intervention

Post-Intervention

Post-Intervention

Total Sample (n=2378)
Allocated to study* (n=103)

Total Sample (n=6038)
Allocated to study* (n=229)

Total Sample (n=995)
Allocated to study* (n=102)

Total Sample (n=3580)
Allocated to study* (n=238)

Analyzed (n=95)
Excluded from analysis:
Chart not found (n=7)
Not meeting inclusion
criteria (n=1)

Analyzed (n=206)
Excluded from analysis:
Chart not found (n=13)
Not meeting inclusion
criteria (n=10)

Analyzed (n=93)
Excluded from analysis:
Chart not found (n=8)
Not meeting inclusion
criteria (n=1)

Analyzed (n=205)
Excluded from analysis:
Chart not found (n=18)
Not meeting inclusion
criteria (n=15)

*Patient charts were allocated to the study using a stratified random sample of patients of equal proportion per physician