

Supplemental Information

Table S1: Antibiotic susceptibilities of the employed bacterial strains against penicillin G, doxycycline, clindamycin, cefazolin, cefuroxime and ceftazidime.

Bacterial strain	Susceptibilities against antibiotics (MIC in µg/ml)					
	Penicillin G	Doxycycline	Clindamycin	Cefazolin	Cefuroxime	Ceftazidime
<i>Streptococcus</i> Group G	<0.064	>2.0	<0.5	<0.064	<0.064	<0.125
<i>Staphylococcus aureus</i> (MSSA)	>0.125	<1.0	<0.25	<4.0	<4.0	>256
<i>Staphylococcus aureus</i> (MRSA)	>256	<1.0	>256	>256	>256	>256
<i>Streptococcus mutans</i>	<0.064	>2.0	<0.5	<0.064	<0.064	<0.125
<i>Streptococcus</i> Group C	<0.064	<1.0	<0.5	<0.064	<0.064	<0.125
<i>Streptococcus pyogenes</i>	<0.064	<1.0	<0.5	<0.064	<0.064	<0.125
<i>Haemophilus influenzae</i>	1.0	<1.0	>4.0	>0.5	<1.0	<0.064
<i>Haemophilus parainfluenzae</i>	>1.0	<1.0	>4.0	>0.5	<1.0	<0.064
<i>Branhamella catarrhalis</i>	>1.0	<1.0	>4.0	>0.5	<1.0	<0.064

MIC = minimal inhibitory concentration