

✓ Mean follow-up score adjusted for pre-intervention score

Hip function after surgery – all English providers

Provider	Outcome (Oxford Hip Score)	95% confidence interval
Adipisicing Consetur	35.87	34.58 to 37.51
Adipisicing Cupidatat	36.04	34.58 to 37.51
Adipisicing Lorem	37.67	35.65 to 39.69
Aliqua Anim	36.59	33.67 to 39.52
Aliqua Cillum	35.09	33.17 to 37.00
Aliqua Dolor	36.43	35.18 to 37.68
Amaboret Deserunt	35.90	34.43 to 37.39
Amentur Consectetur	35.87	34.58 to 37.51

✓ Mean change in score adjusted for pre-intervention score

Improvement in hip function after surgery – all English providers

Provider	Improvement (Oxford Hip Score)	95% confidence interval
Adipisicing Consetur	18.74	15.19 to 22.28
Adipisicing Cupidatat	18.18	16.59 to 19.79
Adipisicing Lorem	20.04	17.84 to 22.25
Aliqua Anim	19.00	15.80 to 22.21
Aliqua Cillum	17.37	15.28 to 19.46
Aliqua Dolor	18.80	17.43 to 20.17
Amaboret Deserunt	18.19	16.57 to 19.81
Amentur Consectetur	18.87	16.58 to 21.17

✓ Proportion reaching a specified threshold adjusted for pre-intervention score

Proportion achieving good hip function – all English providers

Provider	% of patients achieving a good hip function (Oxford Hip Score)	95% confidence interval
Adipisicing Consetur	40	23 to 57
Adipisicing Cupidatat	43	35 to 50
Adipisicing Lorem	49	39 to 60
Aliqua Anim	43	28 to 59
Aliqua Cillum	42	32 to 52
Aliqua Dolor	41	35 to 48
Amaboret Deserunt	37	30 to 45
Amentur Consectetur	47	36 to 58

✓ Proportion reaching a minimally important difference (MID) adjusted for pre-intervention score

Proportion achieving a significant improvement in hip function – all English providers

Provider	% of patients achieving a Minimally Important Difference (OHS)	95% confidence interval
Adipisicing Consetur	84	72 to 96
Adipisicing Cupidatat	82	76 to 87
Adipisicing Lorem	88	81 to 96
Aliqua Anim	89	79 to 100
Aliqua Cillum	79	72 to 86
Aliqua Dolor	85	80 to 89
Amaboret Deserunt	82	76 to 87
Amentur Consectetur	82	74 to 90