

		Percentage of all children treated for fever (95% confidence interval)					
	Yrs since sanctioned CQ use at time of survey	CQ		SP		ACT	
East Africa							
Malawi ^a	13	0.8 (0.6,1.1)		24.9 (23.8,25.9)		0.2 (0.1,0.3)	
Somalia ^a	1	37.0 (29.9,44.0)		14.7 (9.5,19.8)		5.4 (2.1,8.7)	
Tanzania ^{d,‡}	6	0.5 (0.1,0.9)		6.5 (5.0,7.9)		22.2 (19.7,24.7)	
Uganda ^c	0	45.5 (43.7,47.3)		21.4 (19.9,22.9)		3.5 (2.8,4.2)	
Zambia ^d	5	0.6 (0.1,1.1)		28.4 (25.4,31.4)		13.4 (11.1,15.6)	
Zimbabwe ^c	2	51.9 (42.2,61.6)		0.9 (-0.9,2.8)		- (-,-)	
Central Africa							
Angola ^{d,‡}	*	14.7 (9.7,19.7)		0.5 (-0.5,1.5)		4.1 (1.3,6.8)	
Cameroon ^{a,†,‡}	4	8.8 (7.0,10.5)		1.1 (0.5,1.8)		3.0 (2.0,4.1)	
DRC ^d	6	8.5 (7.3,9.7)		3.1 (2.3,3.9)		1.3 (0.8,1.9)	
West Africa							
Benin ^c	2	51.6 (50.0,53.1)		0.5 (0.2,0.7)		0.4 (0.2,0.6)	
Burkina Faso ^a	*	56.8 (53.3,60.3)		0.5 (0.0,1.0)		- (-,-)	
Cote d'Ivoire ^a	3	38.9 (36.6,41.2)		2.0 (1.4,2.7)		2.9 (2.1,3.7)	
Gambia ^a	1	66.9 (62.6,71.2)		14.8 (11.6,18.1)		0.2 (-0.2,0.6)	
Ghana ^a	2	47.4 (43.8,50.9)		0.7 (0.1,1.2)		3.6 (2.2,4.9)	
Guinea-Bissau ^a	-2	64.9 (60.7,69.1)		3.6 (1.9,5.2)		2.6 (1.2,3.9)	
Liberia ^d	3	49.1 (46.4,51.7)		4.4 (3.3,5.4)		9.4 (7.9,11.0)	
Mali ^c	4	28.9 (26.7,31.1)		3.4 (2.5,4.2)		0.2 (0.0,0.5)	
Mauritania ^{b,†}	*	11.7 (9.4,14.0)		6.0 (4.3,7.7)		2.0 (1.0,3.1)	
Niger ^c	1	44.2 (41.6,46.7)		1.0 (0.5,1.5)		- (-,-)	
Senegal ^c	3	10.6 (8.9,12.4)		0.4 (0.1,0.8)		9.7 (8.1,11.4)	
Togo ^a	*	36.7 (33.1,40.3)		3.8 (2.4,5.3)		1.6 (0.7,2.6)	