

Database: MEDLINE <1966 to July 2003>

Search Strategy

- 1 exp gastroenteritis/
- 2 gastroenteritis.tw.
- 3 exp diarrhea/
- 4 or/1-3
- 5 exp fluid therapy/
- 6 exp water-electrolyte balance/
- 7 exp water-electrolyte imbalance/
- 8 exp rehydration solutions/
- 9 rehydrat\$.mp.
- 10 exp infusions, intravenous/
- 11 or/5-10
- 12 exp Feeding methods/
- 13 nose feed\$.mp.
- 14 nasogastric feed\$.mp.
- 15 naso-gastric feed\$.mp.
- 16 ng feed\$.mp.
- 17 ng tube.mp.
- 18 "NASOGASTRIC TUBE".mp.
- 19 or/12-18
- 20 11 or 19
- 21 4 and 20
- 22 limit 21 to (all infant < birth to 23 months > or "all child < 0 to 18 years >" or newborn infant < birth to 1 month > or infant < 1 to 23 months > or preschool child < 2 to 5 years > or child < 6 to 12 years > or adolescence < 13 to 18 years >)
- 23 random\$.mp.
- 24 multicenter study.mp.
- 25 exp meta-analysis/
- 26 or/23-25
- 27 22 and 26

Database: CENTRAL <2003, Issue 2>

Search Strategy

1. gastroenteritis and oral rehydration therapy
2. oral rehydration therapy and intravenous
3. gastroenteritis and intravenous therapy
4. oral rehydration, dehydration and diarrhea

Database: EMBASE <1980 to July 2003>

Search Strategy

- 1 exp acute gastroenteritis/
- 2 exp gastroenteritis/
- 3 exp infantile gastroenteritis/
- 4 exp diarrhea/
- 5 or/1-4
- 6 exp rehydration/
- 7 exp oral rehydration solution/
- 8 oral rehydration.tw.
- 9 exp intravenous feeding/
- 10 exp fluid therapy/
- 11 exp intravenous administration/
- 12 or/6-11
- 13 Nose feeding/
- 14 nasogastric feed\$.mp.
- 15 naso-gastric.mp.
- 16 nose feed\$.mp.
- 17 ng feed\$.mp.
- 18 ng tube.mp.
- 19 nasogastric tube.mp.
- 20 or/13-19
- 21 12 or 20
- 22 21 and 5
- 23 limit 22 to (human and ("all child < 0 to 18 years >" or newborn infant < birth to 1 month > or infant < 1 to 23 months > or preschool child < 2 to 5 years > or child < 6 to 12 years > or adolescence < 13 to 18 years >))
- 24 exp meta analysis/
- 25 exp multicenter study/
- 26 multicenter study.mp.
- 27 random\$.mp.
- 28 or/24-27
- 29 23 and 28