

Additional Table S1 Characteristics of the 452 primary breast tumors, and relation to metastasis-free survival.

	Number of patients	5-y MFS	P-value ^a
<i>Total</i>	452	73.4 %	-
<i>Age</i>			
≤50	96	83.3 %	NS
>50	356	80.5 %	
<i>SBR histological grade^{b,c}</i>			
I	55	92.5 %	0.000064^f
II	228	75.3 %	
III	159	64.3 %	
<i>Lymph node status^d</i>			
0	115	79.0 %	0.00000026^f
1-3	237	76.8 %	
>3	99	58.6 %	
<i>Macroscopic tumor size^e</i>			
≤25mm	217	82.7 %	0.000011
>25mm	228	64.1 %	
<i>ERα</i>			
Negative	117	60.1 %	0.0046
Positive	335	78.1 %	
<i>PR</i>			
Negative	194	62.6 %	0.0007
positive	258	81.4 %	
<i>ERBB2</i>			
Negative	351	75.3 %	NS
Positive	101	66.8 %	
<i>HR/ERBB2 subgroups</i>			
HR+ ERBB2+	53	79.0 %	0.0049^f
HR + ERBB2-	287	78.3 %	
HR - ERBB2+	48	53.4 %	
HR - ERBB2 -	64	62.0 %	
<i>Histology</i>			
Ductal	388	73.5 %	NS ^f
Lobular	29	79.3 %	
Others	35	68.3 %	

^aLog-rank test. NS: not significant.

^bScarff Bloom Richardson classification.

^cInformation available for 442 patients.

^dInformation available for 451 patients.

^eInformation available for 445 patients.

^fGlobal comparison of all subgroups of a category.