

Supplementary Table 3 - Multiple medication overuse headache sub-diagnoses and specific combination of medication overuse

Combination of MOH diagnoses	Denmark	Germany	Italy	Spain	Argentina	Chile	Total
N	125	101	117	92	126	108	669
Single MOH type	74.4	78.2	93.2	87.0	81.7	38.9	75.6
Two MOH types	24.8	17.8	6.8	13.0	17.5	57.4	22.9
Three MOH types	0.8	4.0	0.0	0.0	0.8	3.7	1.5
Triptans and ergotamines	0.0	0.0	0.0	2.2	1.6	0.9	0.7
Triptans and simple analgesics	5.6	9.9	2.6	4.3	0.8	0.9	3.9
Triptans and combination-drugs	4.0	1.0	1.7	0.0	0.0	0.9	1.3
Triptans and poly-overuse	1.6	0.0	0.0	0.0	0.0	0.0	0.3
Ergotamines and simple analgesics	0.8	0.0	0.9	3.3	14.3	1.9	3.7
Ergotamines and combination-drugs	0.8	0.0	0.0	1.1	0.8	45.4	7.8
Simple analgesics and opioids	2.4	2.0	0.0	0.0	0.0	0.0	0.7
Simple analgesics and combination-drugs	8.8	3.0	1.7	2.2	0.0	6.5	3.7
Simple analgesics and poly-overuse	0.8	0.0	0.0	0.0	0.0	0.0	0.1
Opioids and combination-drugs	0.0	1.0	0.0	0.0	0.0	0.0	0.1
Combination-drugs and poly-overuse	0.0	1.0	0.0	0.0	0.0	0.0	0.3
Ergotamines, simple analgesics and combination-drugs	0.0	0.0	0.0	0.0	0.0	2.8	0.4
Triptans, simple analgesics and combination-drugs	0.8	1.0	0.0	0.0	0.0	0.9	0.4
Simple analgesics, opioids and combination-drugs	0.0	1,0	0.0	0.0	0.0	0.0	0.1
Triptans, simple analgesics and opioids	0.0	1,0	0.0	0.0	0.0	0.0	0.1
Ergotamines, triptans and simple analgesics	0.0	1,0	0.0	0.0	0.8	0.0	0.3

The table characterizes the proportion of patients with multiple medication overuse headache (MOH) sub-diagnoses and the specific medication overuse profile. The data are shown as percentages.

