

Additional file 1

Assessment of pesticides in surface water samples from Swedish agricultural areas by integrated bioanalysis and chemical analysis

Johan Lundqvist^{*1}, Claudia von Brömssen², Anna Kjerstine Rosenmai¹, Åsa Ohlsson¹, Theo Le Godec¹, Ove Jonsson³, Jenny Kreuger³ and Agneta Oskarsson¹

¹Department of Biomedicine and Veterinary Public Health, ²Department of Energy and Technology, ³Department of Aquatic Sciences and Assessment
Swedish University of Agricultural Sciences, Box 7028, SE-750 07 Uppsala, Sweden.

Table of contents

Table S1	List of analyzed pesticides, CAS number and analytical method used for each compound
Table S2	Substances excluded from the statistical analysis, based on few observations (less than four observations).
Table S3	Recovery for pesticides normally analyzed with an LC-MS/MS method, when extracted according to the GC-MS method
Table S4	Bioactivity and cell viability (fold change) and pesticide concentrations ($\mu\text{g/l}$) in the water samples
Figure S1	Cell viability of H295R, HepG2, and VM7Luc4E2 cells, after exposure to concentrated water samples for 24 h

Figure S1. Cell viability of H295R, HepG2, and VM7Luc4E2 cells, after exposure to concentrated water samples for 24 h (mean, n=4). The solid line indicates the threshold for perceived compromised cell viability of 75%. For standard deviation, refer to Table S4.

