

Table of contents

Additional table 1: Genes primers sequences used for RT-qPCR	1
Additional figure 1: Venn diagram showing the number of samples shared between the different analysis techniques that were used. The blue circle represents the number of samples with results obtained from transcriptomics. The green circle represents the number of samples with results for both metabolomics and microbial composition. The red circle represents samples with RT-qPCR results.	2
Additional figure 2: On 543 children, the percentage of preterm infants with high ASQ (> 185) in blue and low ASQ (<185) in red; $P = 3.756e-07$ with G test. Enterotype 1 (n=236), enterotype 2 (n=68), enterotype 3 (n=60), enterotype 4 (n=62), enterotype 5 (n=50) and enterotype 6 (n=67).....	3
Additional figure 3: Frequency of preterm infants with high ASQ (> 185) in blue and low ASQ (<185) in red depending on weight at birth. $P = 0.1147$ with G test. NBW : Normal birth weight (n=32), LBW : Low birth weight (n=17) and VLBW : Very low birth weight (n=24).	4
Additional figure 4: Relative abundance of 7 major TCA cycle metabolites by clusters: citrate (A), aconitate (B), isocitrate (C), alpha ketoglutarate (D), succinate (E), fumarate (F), malate (G). Dunn test. ** = $P < 0.01$; *** = $P < 0.001$, ns: non-significant. Cluster A (n=20), Cluster B (n=20), Cluster C (n=20).....	5
Additional figure 5: Relative abundance of 6 major TCA cycle metabolites by ASQ: citrate (A), aconitate (B), alpha ketoglutarate (C), succinate (D), fumarate (E), malate (F). ASQ <185 (n=15), ASQ >185 (n=45).	6

Additional table 1: **Genes primers sequences used for RT-qPCR**


Gene	Forward	Reverse	Ref
S100A8	GCTAGAGACCGAGTGTCTCCTCAG	GCCCATCTTTATCACCAGAATG	1
S100A9	TGGAGGACCTGGACACAAATG	TCGTCACCCTCGTGCATCTT	1
CAMP/LL37	GACACAGCAGTCACAGAGGAT	TCACAACTGATGTCAAAGGAGCC	2
NAMPT	AGGGTTACAAGTTGCTGCCACC	CTCCACCAGAACCGAAGGCAAT	3

1 Yan LL, Huang YJ, Yi X, et al. Effects of silencing S100A8 and S100A9 with small interfering RNA on the migration of CNE1 nasopharyngeal carcinoma cells. *Oncol Lett* 2015; 9: 2534.


2 Cathelicidin (CAMP) Human qPCR Primer Pair (NM_004345) – HP207673 | OriGene. https://www.origene.com/catalog/gene-expression/qpcr-primer-pairs/hp207673/cathelicidin-camp-human-qpcr-primer-pair-nm_004345.

3 Visfatin (NAMPT) Human qPCR Primer Pair (NM_005746) – HP208829 | OriGene. https://www.origene.com/catalog/gene-expression/qpcr-primer-pairs/hp208829/visfatin-nampt-human-qpcr-primer-pair-nm_005746.


Additional figure 1: **Venn diagram showing the number of samples shared between the different analysis techniques that were used.** The blue circle represents the number of samples with results obtained from transcriptomics. The green circle represents the number of samples with results for both metabolomics and microbial composition. The red circle represents samples with RT-qPCR results.




Additional figure 2: **On 543 children, the percentage of preterm infants with high ASQ (> 185) in blue and low ASQ (<185) in red; $P = 3.756e-07$ with G test. Enterotype 1 (n=236), enterotype 2 (n=68), enterotype 3 (n=60), enterotype 4 (n=62), enterotype 5 (n=50) and enterotype 6 (n=67).**



Additional figure 3: **Frequency of preterm infants with high ASQ (> 185) in blue and low ASQ (<185) in red depending on weight at birth.** P = 0.1147 with G test. NBW : Normal birth weight (n=32), LBW : Low birth weight (n=17) and VLBW : Very low birth weight (n=24).



Additional figure 4: **Relative abundance of 7 major TCA cycle metabolites by clusters:** citrate (A), aconitate (B), isocitrate (C), alpha ketoglutarate (D), succinate (E), fumarate (F), malate (G). Dunn test. ** = $P < 0.01$; *** = $P < 0.001$, ns: non-significant. Cluster A (n=20), Cluster B (n=20), Cluster C (n=20).



Additional figure 5: **Relative abundance of 6 major TCA cycle metabolites by ASQ**: citrate (A), aconitate (B), alpha ketoglutarate (C), succinate (D), fumarate (E), malate (F). ASQ <185 (n=15), ASQ >185 (n=45).

