

Table S1 Information alert form for trauma patients

Age	☐ 0-3 years old ☐ 3-14 years old ☐ 14-60 years old ☐ over 60 years old
Time	____(year) / ____ (month) / ____ (date) / ____ (h) / ____ (min)
Mechanism of injury	☐ Traffic accident ☐ Fall from a height ☐ Burns /scalds / explosion injuries ☐ Stab wound ☐ Others
Injuries	☐ Head and neck ☐ Face ☐ Chest ☐ Abdomen ☐ Limbs /pelvis /buttocks ☐ Body surfaces Other: _____
Signs	Blood pressure: ____/____mmHg Heart rate: ____/min Respiratory rate: ____/min SpO ₂ : ____% Consciousness: ☐ Comatose ☐ Alert
Treatment	☐ Cervical collar immobilization ☐ Oxygen therapy/Oral cavity clearance ☐ Dressing and hemostasis ☐ Splinting ☐ Pelvic binder fixation Other: _____
Others	

The criteria for initiating a multidisciplinary trauma team are as follows: 1) Trauma-induced cardiac arrest requiring cardiopulmonary resuscitation, Glasgow Coma Scale (GCS) ≤ 8 , systolic blood pressure < 90 mmHg (1 mmHg = 0.133 kPa), heart rate > 120 beats/min, or hypothermia. 2) Post-traumatic respiratory distress or the need for endotracheal intubation. 3) Penetrating injuries to the head, neck, or torso. 4) Hemodynamically unstable pelvic fractures, limb mutilation, or severe combined injuries.