

Extent of disease affects the usefulness of fecal biomarkers in ulcerative colitis

Authors:

Akihito Sakuraba¹, Nobuki Nemoto¹, Noritaka Hibi¹, Ryo Ozaki¹, Sotaro Tokunaga¹, Oki Kikuchi¹, Shintaro Minowa¹, Tatsuya Mitsui¹, Miki Miura¹, Daisuke Saito¹, Mari Hayashida¹, Jun Miyoshi¹, Minoru Matsuura¹, Masayoshi Yoneyama², Hiroaki Ohnishi³, Tadakazu Hisamatsu^{1*}

Affiliation:

- 1) Department of Gastroenterology and Hepatology, Kyorin University School of Medicine, Tokyo, Japan
- 2) Department of Clinical Laboratory, Kyorin University Hospital, Tokyo, Japan.
- 3) Department of Laboratory Medicine, Kyorin University School of Medicine, Tokyo, Japan.

*** Corresponding author:**

Tadakazu Hisamatsu, M.D., Ph.D., FACP, AGAF
Department of Gastroenterology and Hepatology,
Kyorin University School of Medicine.
6-20-2 Shinkawa, Mitaka-shi, Tokyo 181-8611, Japan

Tel.: +81-422-47-5511 (ext. 5279)

Fax: +81-422-44-0655

E-mail: thisamatsu@ks.kyorin-u.ac.jp

Running title: Disease extent and fecal biomarkers in UC

Supplemental Table 1. Mayo Endoscopic Subscore (MES)

Score	Disease Activity	Endoscopic Features (Descriptors)
0	Normal or inactive	None
1	Mild	Erythema, decreased vascular pattern, mild friability
2	Moderate	Marked erythema, absent vascular pattern, friability, erosions
3	Sever	Spontaneous bleeding, ulceration