

Supplementary Table 3. Reclassification and discrimination statistics for outcomes.

Variables	C statistic		IDI		NRI	
	Estimate	<i>P</i> value	Estimate (95% CI)	<i>P</i> value	Estimate (95% CI)	<i>P</i> value
28-day mortality						
Conventional model	0.754		Reference		Reference	
Conventional model + ALBI	0.761	0.0399	0.011 (0.006-0.016)	<0.001	0.242 (0.148-0.335)	<0.001
Conventional model + SOFA	0.758	0.1135	0.004 (0.001-0.007)	0.02	0.117 (0.023-0.211)	0.02
Conventional model + MELD	0.804	<0.001	0.08 (0.06-0.09)	<0.001	0.603 (0.513-0.693)	<0.001
90-day mortality						
Conventional model	0.746		Reference		Reference	
Conventional model + ALBI	0.753	0.02	0.009 (0.005-0.013)	<0.001	0.202 (0.114-0.290)	0.003
Conventional model + SOFA	0.748	0.21	0.004 (0.001-0.006)	0.01	0.114 (0.027-0.202)	0.01
Conventional model + MELD	0.799	<0.001	0.08 (0.07-0.09)	<0.001	0.618 (0.534-0.702)	<0.001

Conventional model: age, gender, ethnicity, white blood cell, hypertension, heart failure, respiratory failure, diabetes, vasopressin, continuous renal replacement therapy.