Additional file 4. Table univariate analyses for associations between patient characteristics and willingness

	Willingness if GP proposes (n = 196)			Willingness if specialist proposes (n = 183)			Willingness if pharmacist proposes (n = 195)		
	p value	OR	95 % CI	p value	OR	95 % CI	p value	OR	95 % CI
Age group[a]									
65-69	0.98	0.99	0.45-2.19	0.68	1.19	0.53-2.69	0.34	1.44	0.68-3.06
70-74	0.77	0.88	0.38-2.05	0.78	0.88	0.37-2.09	0.65	1.21	0.54-2.70
> 75	0.85	1.10	0.40-3.06	0.70	1.23	0.44-3.46	0.77	0.86	0.32-2.31
Sex (male)	0.48	0.78	0.38-1.57	0.97	1.02	0.50-2.04	0.79	0.91	0.48-1.74
Number of medicines	0.79	1.03	0.84-1.27	0.82	0.98	0.80-1.20	**0.16**	**0.86**	**0.71-1.06**
Education level[c]									
Junior high school	0.26	0.58	0.23-1.48	**0.15**	**0.50**	**0.19-1.27**	**0.01**	**0.27**	**0.11-0.69**
Senior high school	0.45	0.73	0.33-1.64	0.69	0.85	0.37-1.94	**0.03**	**0.43**	**0.21-0.90**
University degree	0.29	0.62	0.26-1.49	0.23	0.58	0.24-1.40	**0.01**	**0.35**	**0.15-0.80**
Type of medicines									
Using >1 glucose-lowering medicine[c]	**0.04**	**0.53**	**0.29-0.98**	**0.13**	**0.62**	**0.33-1.15**	0.40	0.78	0.44-1.38
Using blood pressure-lowering medicines[d]	0.43	1.29	0.68-2.45	0.66	1.16	0.60-2.23	0.88	0.96	0.52-1.76
Using lipid-lowering medicines[e]	**0.08**	**1.91**	**0.92-3.95**	0.29	1.47	0.72-2.99	0.82	1.08	0.57-2.03
Using CAM[f]	0.76	1.11	0.58-2.12	0.49	1.28	0.64-2.54	0.74	1.11	0.60-2.03

Abbreviations: GP: general practitioner, OR: Odds ratio, CI: confidence interval, CAM: complementary and alternative medicines

[a] Age group 60-64 years as reference

[b] Primary school or no school as reference

[c] Using 1 glucose-lowering medicines as reference

[d] Using no blood pressure-lowering medicines as reference

[e] Using no lipid-lowering medicines as reference

[f] Using no CAM as reference