

Supplementary Table S6. Characteristics for females and males.

a) Characteristics	All Participants		GM-PCP Subgroup		
	Females (n = 160)	Males (n = 87)	Physician Type	Females (n = 59)	Males (n = 37)
Age (y, mean ± sd)	81.5 ± 7.0	82.0 ± 7.5	GM	83.0 ± 6.0	84.7 ± 7.16
			PCP	83.0 ± 5.9	84.7 ± 7.14
Sex (women, %)	100.0	0.0	GM	100.0	0.0
			PCP	100.0	0.0
Cognitive Impairments (%)	22.5	32.2	GM	40.7	51.4
			PCP	39.7	35.1
NCI (%)	77.5	67.8	GM	59.3	48.7
			PCP	60.3	64.9
MCI (%)	18.8	28.7	GM	28.8	35.1
			PCP	29.3	35.1
Dementia (%)	3.7	3.5	GM	11.9	16.2
			PCP	10.4	0.0
Education (y, mean ± sd)	12.9 ± 3.3	13.0 ± 4.6	GM	13.8 ± 3.3	14.1 ± 4.34
			PCP	13.8 ± 3.4	14.1 ± 4.50
Currently Married (%)	42.8	69.8	GM	50.8	73.0
			PCP	50.8	73.0
Lives Alone (%)	50.9	29.9	GM	34.5	21.6
			PCP	45.8	27.0
Ever Smoked (%)	48.8	65.1	GM	54.2	62.2
			PCP	64.4	67.6
Hypertension (%)	64.5	59.7	GM	54.7	61.1
			PCP	66.7	64.9
Diabetes (%)	22.0	34.2	GM	15.1	16.7
			PCP	16.7	18.9
Stroke/TIA (%)	7.3	7.9	GM	13.2	8.3
			PCP	13.0	10.8
Depression (%)	20.9	19.5	GM	20.3	8.1
			PCP	22.4	18.9
Mini-Cog (mean ± sd)	3.9 ± 1.4	3.3 ± 1.5	GM	3.9 ± 1.3	3.5 ± 1.5
			PCP	3.7 ± 1.4	3.3 ± 1.4

MoCA (mean ± sd)	20.6 ± 5.5	21.1 ± 5.4	GM	21.6 ± 5.5	22.0 ± 4.9
			PCP	21.5 ± 5.6	21.9 ± 4.5
FAST (mean ± sd)	2.4 ± 1.4	2.7 ± 1.4	GM	2.5 ± 1.2	3.1 ± 1.4
			PCP	2.4 ± 1.2	2.5 ± 1.1
Medications >7 (%)	36.1	54.0	GM	32.2	43.2
			PCP	40.4	43.2
Number of Problems (mean ± sd)	7.6 ± 4.6	8.3 ± 3.4	GM	8.1 ± 3.5	8.1 ± 2.4
			PCP	7.1 ± 4.2	7.9 ± 3.0
CFS (mean ± sd)	3.7 ± 1.2	3.9 ± 1.2	GM	3.9 ± 1.4	4.2 ± 1.2
			PCP	3.7 ± 0.9	4.1 ± 0.9
eFI-CGA (mean ± sd)	0.27 ± 0.13	0.29 ± 0.11	GM	0.28 ± 0.13	0.31 ± 0.12
			PCP	0.27 ± 0.12	0.30 ± 0.10
NCI eFI-CGA (mean ± sd)	0.25 ± 0.13	0.28 ± 0.11	GM	0.23 ± 0.11	0.25 ± 0.09
			PCP	0.23 ± 0.10	0.28 ± 0.09
MCI eFI-CGA (mean ± sd)	0.33 ± 0.12	0.32 ± 0.10	GM	0.35 ± 0.12	0.36 ± 0.12
			PCP	0.32 ± 0.14	0.33 ± 0.10
Dementia eFI-CGA (mean ± sd)	0.33 ± 0.11	0.38 ± 0.10	GM	0.35 ± 0.14	0.38 ± 0.10
			PCP	0.35 ± 0.12	N/A