

Supplementary Table S7. Multivariable analysis of factors associated with hematological adverse events using full model without p-value screening (n=311)

Variable	aPR	95% CI	p-value
Residence			
Urban	1.00 (Ref)		
Rural	1.34	1.09 – 1.64	0.005
Nature of Employment			
Employed	1.00 (Ref)		
Self-employed	1.13	0.84 – 1.52	0.411
Unemployed	0.75	0.56 – 1.02	0.071
Marital Status			
Single	1.00 (Ref)		
Married	0.66	0.53 – 0.84	0.001
Divorced/Widowed	0.91	0.63 – 1.31	0.624
Cigarette Smoking			
No	1.00 (Ref)		
Yes	1.67	1.35 – 2.07	<0.001
HIV Status			
Negative	1.00 (Ref)		
Positive	1.16	0.92 – 1.45	0.207
Alcohol Use			
No	1.00 (Ref)		

Yes	0.84	0.69 – 1.02	0.084
Previous History of TB Treatment			
No	1.00 (Ref)		
Yes	1.04	0.85 – 1.27	0.708

Note: Model adjusted for all a priori selected covariates shown. Modified Poisson regression with robust standard errors. Listwise deletion resulted in a sample size of 311 patients due to missing covariate data.